

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	Kidney Transplant
Policy Number	7.02.04
Category	Transplants
Original Effective Date	04/19/00
Committee Approval Date	08/16/01, 06/20/02, 05/21/03, 02/19/04, 02/17/05, 01/19/06, 02/15/07, 01/17/08, 03/19/09, 03/18/10, 03/17/11, 02/16/12, 01/17/13, 02/20/14, 03/24/22
Current Effective Date	03/24/22
Archived Date	02/19/15
Archive Review Date	03/17/16, 03/16/17, 03/15/18, 03/21/19, 3/19/20, 03/18/21, 03/24/22
Product Disclaimer	<ul style="list-style-type: none"> If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. If a commercial product (including an Essential Plan or Child Health Plus product), medical policy criteria apply to the benefit. If a Medicaid product covers a specific service, and there are no New York State guidelines (eMedNY) criteria, medical policy criteria apply to the benefit. If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

- I. Based upon our criteria and assessment of the peer-reviewed literature, kidney transplants for carefully selected candidates, who have documentation of progressive end-stage renal disease (ESRD) and no immediate life-threatening conditions, have been medically proven to be effective and, therefore, are considered **medically appropriate** for patients with:
 - A. A measured (actual urinary collection) creatinine clearance level or calculated GFR (Cockcroft-Gault) or other reliable formula) less than or equal to 20ml/min; or
 - B. The initiation of dialysis.
- II. Recipient Selection
 - A. Each individual considered for renal transplantation will have an evaluation completed by the transplant center for potential difficulties that would complicate and diminish the success of transplantation. Consideration will be given to the patient's risk of death without transplantation, along with the presence and severity of potential contraindications to transplantation. Candidates considered for transplant must be psychologically stable, demonstrate motivation and compliance, and have no ongoing problems with drug or alcohol abuse.
 - B. Conditions that preclude proceeding to transplantation include, but are not limited to:
 1. Metastatic cancer;
 2. Presence of malignancy (other than non-melanoma skin cancers), unless malignancy has been completely resected, or unless (upon medical review) it is determined that malignancy has been treated with small likelihood of recurrence and acceptable future risks;
 3. Ongoing or recurring infections that are not effectively treated;
 4. Serious cardiac or other ongoing insufficiencies that create an inability to tolerate transplant surgery; and
 5. Demonstrated non-compliance, which places the organ at risk by not adhering to medical recommendations.
 - C. Renal transplantation in the context of asymptomatic HIV infection is rapidly evolving in the setting of highly active antiretroviral therapy (HAART). Currently, the United Network for Organ Sharing (UNOS) states that asymptomatic HIV-positive patients should not necessarily be excluded for candidacy for organ transplantation. In 2001, the Clinical Practice Committee of the American Society of Transplantation proposed that the presence of

Medical Policy: KIDNEY TRANSPLANT

Policy Number: 7.02.04

Page: 2 of 6

AIDS could be considered a contraindication to kidney transplant, unless the patient meets **ALL** of the following criteria:

1. CD4 count greater than 200 cells/mm³.
2. Undetectable HIV-1RNA.
3. On stable anti-retroviral therapy for greater than three months.
4. No other complications from AIDS (e.g., opportunistic infection, including aspergillus, tuberculosis, coccidioidomycosis; resistant fungal infections, Kaposi's sarcoma, or other neoplasm); and
5. All other criteria for transplantation.

- D. Diabetic complications often fall into the realm of relative contraindications (except for significant cardiovascular disease); however, renal transplantation is associated with improved survival in patients with ESRD caused by type 1 diabetes mellitus. Patients with diabetes may be candidates for combined kidney-pancreas transplantation.

POLICY GUIDELINES

- I. Prior authorization requirements are contract dependent. Approvals for all transplants, including arrangements with an approved transplant center, may be required.
- II. Pre-transplant evaluation documentation must include the following clinical information (if testing is unable to be performed, the rationale for not performing the testing should be included in the documentation):
 - A. Clinical Evaluation:
 1. Confirmation of diagnosis;
 2. Identification of comorbidities;
 3. Treatment of co-morbidities;
 4. Current assessment of co-morbidities; and
 5. Consult notes (if applicable).
 - B. Psycho-Social Evaluation:
 1. Karnofsky performance score; and
 2. Identification of stressors (family support, noncompliance issues, motivational issues, alcohol, or substance abuse).
 - C. Dental Evaluation.
 - D. Lab Tests:
 1. CBC, metabolic profile;
 2. Serologies: CMV;
 3. Hepatitis B and C; and
 4. HIV Testing.
 - E. Cardiac Assessment:
 1. 12 Lead EKG; and
 2. Stress echo or MUGA Scan.
 - F. Pulmonary Assessment:
 1. Chest x-ray;
 2. Pulmonary function tests (PFTs); and
 3. Low-dose screening CT for individuals considered high-risk for lung cancer (e.g., 20- to 30-pack history of smoking).
 - G. Age-appropriate Screening Tests:
 1. Age 45 years or greater:
 - a. Colonoscopy (within 10 years); or
 - b. Flexible sigmoidoscopy (within five years); or

Medical Policy: KIDNEY TRANSPLANT

Policy Number: 7.02.04

Page: 3 of 6

- c. Guaiac stool testing (within one year); or
 - d. Rationale of contraindication to testing (if applicable).
 2. Age 21 to 70 years, as appropriate:
 - a. Pap smear (within three years).
 3. Age 40 years or greater, as appropriate:
 - a. Mammogram (within two years).
- III. Hepatitis C virus (HCV) infections are common among patient with chronic renal failure and result in significant morbidity and mortality. Therefore, the assessment of hepatitis C virus infection in the potential recipient has a major clinical significance. HCV infections are associated with an increased risk of death, irrespective of whether of the patient stays on dialysis or has a renal transplant. Transplantation has a beneficial, rather than adverse, effect on long-term survival in an anti-HCV-positive patient. An anti-HCV-positive status is not a contradiction for renal transplantation.
- IV. Living Donation: Any person who gives consent to be a live organ donor should be competent, willing to donate, free from coercion, medically and psychologically suitable, fully informed of the risks and benefits as a donor, and fully informed of the risks, benefits, and alternative treatment available to the recipient. The benefits to both donor and recipient must outweigh the risks associated with the donation and transplantation for the living donor organ.
- V. Candidates may be wait-listed at more than one transplant center. Since waiting time priority is first calculated among candidates at all hospitals within the local donation area, listing at transplant centers in different local allocation areas is recommended. Requirements for multiple-listed candidates may vary among transplant centers. When possible, results of tests used in the evaluation for the transplant at one center should be used at subsequent centers where the patient is listed.

DESCRIPTION

A kidney transplant involves the surgical removal of a kidney from a cadaver donor or a matched living donor (related or unrelated) into a recipient for treatment of ESRD.

In the last 10 to 15 years, the incidence of ESRD in patients aged 65 years and older has doubled. Renal transplantation confers substantial survival advantages over dialysis for patients greater than age 60 years. During the first year of renal replacement treatment, transplant and dialysis have similar survival rates. However, from the second-year onward transplantation has demonstrated significant benefit over dialysis in the older patient with ESRD who meets the other criteria for transplant. The beneficial effects of transplantation over dialysis begin to disappear when the average age exceeds 65 years.

Re-transplantation is far more common in kidney transplantation than in other solid organ transplants.

RATIONALE

Kidney transplant is an established treatment option for patients with progressive or end-stage renal disease. Good outcomes have been achieved outside the investigational setting.

Solid organ transplantation for candidates who are HIV-positive has long been controversial, due to the long-term prognosis for HIV positivity, and the impact of immunosuppression on HIV disease. Although HIV-positive transplant recipients may be a research interest of some transplant centers, the minimal data regarding long-term outcome in these patients consist primarily of case reports and abstract presentations of liver and kidney recipients. Nevertheless, some transplant surgeons argue that HIV positivity is no longer an absolute contraindication to transplant, due to the advent of HAART, which has markedly changed the natural history of the disease. Furthermore, UNOS has indicated that asymptomatic HIV-positive patients should not necessarily be excluded from candidacy for organ transplantation, stating, "A potential candidate for organ transplantation whose test for HIV is positive but who is in an asymptomatic state should not necessarily be excluded from candidacy for organ transplantation, but should be advised that he or she may be at increased risk of morbidity and mortality because of immunosuppressive therapy." In 2001, the Clinical Practice

Medical Policy: KIDNEY TRANSPLANT**Policy Number: 7.02.04****Page: 4 of 6**

Committee of the American Society of Transplantation proposed that the presence of AIDS could be considered a contraindication to kidney transplant, unless the specific criteria were present (*refer to Policy Statement IIC*).

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- ***CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.***
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

CPT Codes

Code	Description
50300	Donor nephrectomy (including cold preservation); from cadaver donor, unilateral or bilateral
50320	Donor nephrectomy (including cold preservation); open, from living donor
50323	Backbench standard preparation of cadaver donor renal allograft prior to transplantation, including dissection and removal of perinephric fat, diaphragmatic and retroperitoneal attachments, excision of adrenal gland, and preparation of ureter(s), renal vein(s), and renal artery(s), ligating branches, as necessary
50325	Backbench standard preparation of living donor renal allograft (open or laparoscopic) prior to transplantation, including dissection and removal of perinephric fat and preparation of ureter(s), renal vein(s), and renal artery(s), ligating branches, as necessary
50327	Backbench reconstruction of cadaver or living donor renal allograft prior to transplantation; venous anastomosis, each
50340	Recipient nephrectomy (separate procedure)
50360	Renal allotransplantation, implantation of graft; without recipient nephrectomy
50365	Renal allotransplantation, implantation of graft; with recipient nephrectomy
50370	Removal of transplanted renal allograft
50380	Renal autotransplantation, reimplantation of kidney

Copyright © 2022 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
No codes	

ICD10 Codes

Code	Description
N18.1-N18.9	Chronic kidney disease (CKD) (code range)

REFERENCES

American Society of Transplant Surgeons (ASTS), The American Society of Transplantation (AST), The Association of Organ Procurement Organizations (AOPO), et al. Statement on transplantation of organs from HIV-infected deceased

Medical Policy: KIDNEY TRANSPLANT

Policy Number: 7.02.04

Page: 5 of 6

donors. 2011; https://asts.org/docs/default-source/position-statements/transplantation-of-organs-from-hiv-infected-deceased-donors-july-22-2011.pdf?sfvrsn=fbae5a20_4. Accessed 02/03/2022.

*Andrews PA. Renal Transplantation. BMJ 2002 Mar; 324:530-4.

*Braun WE. Update on kidney transplantation: increasing clinical success, expanding waiting lists. Cleveland Clin J Med 2002 Jun;69(6):501-4.

Choi A, Han JJ. First porcine to human kidney transplantation performed in the United States. Artif Organs. 2022 Feb;46(2):178-179.

*Dominguez-Gil B, et al. Transplantation in the patient with hepatitis C. Transpl Int 2009;22(12):1117-31.

Englum BA, et al. Outcomes in kidney transplant recipients from older living donors. Transplantation 2015 Feb;99(2):309-15.

*Frassetto LA, et al. Renal transplantation in patients with HIV. Nat. Rev Nephrol 2009;5:582–9.

*Gruber SA, et al. Preliminary experience with renal transplantation in HIV-recipients: low acute rejection and infection rates. Transplant 2008 Jul;86(2):269-74.

Gusukuma LW, et al. Outcomes in obese kidney transplant recipients. Transplant Proc 2014 Dec;46(10):3416-9.

Ingsathit A, et al. Long-term outcome of kidney retransplantation in comparison with first kidney transplantation: a report from the Thai Transplantation Registry. Transplant Proc 2013 May;45(4):1427-30.

*Izquierdo L, et al. Third and fourth kidney transplant: still a reasonable option. Transplant Proc 2010;42(7):2498–502.

Kutner NG, et al. Perspectives on the new kidney disease education benefit: early awareness, race and kidney transplant access in ta USRDS study. Am J Transplant 2012 Apr;12(4):1017-23.

Malat GE, et al. High frequency of rejections in HIV-positive recipients of kidney transplantation: a single center prospective trial. Transplantation 2012 Nov 27;94(10):1020-4.

*Martina MN, et al. Kidney transplantation and waiting list for renal transplantation for human immunodeficiency virus patients. Transplant Proc 2011 Jul-Aug;43(6):2179-89.

*Mazuecos A, et al. HIV infection and renal transplantation. Nephrol Dial Transplant 2011;26(4):1401–7.

Mgbako O, et al. Allowing HIV-positive organ donation: ethical, legal and operational considerations. Am J Transplan 2013 Jul;13(7):1636-42.

Muller E, et al. Renal transplantation between HIV-positive donors and recipients justified. S Afr Med J 2012 Mar 2;102(6):497-8.

Nicoletto BB, et al. Effects of obesity on kidney transplantation outcomes: a systematic review and meta-analysis. Transplantation 2014 Jul 27;98(21):167-76.

Porrett PM, et al. First clinical-grade porcine kidney xenotransplant using a human decedent model. Am J Transplant. 2022 Jan 20.

*Steinman TI, et al. The clinical practice committee of the American Society of Transplantation. Guidelines for the referral and management of patients eligible for solid organ transplantation. Transplant 2001;71(9):1189-204.

*Stock PG, et al. Outcomes of kidney transplantation in HIV-infected recipients. N Engl J Med 2010;363:2004-14.

*Trullas JC, et al. Renal transplantation in HIV-infected patients: 2010 update. Kidney Int 2011; 79(8):825-42.

Van Arendonk KJ, et al. Practice patterns and outcomes in retransplantation among pediatric kidney transplant recipients. Transplantation 2013 Jun 15;95(11):1360-8.

Medical Policy: KIDNEY TRANSPLANT

Policy Number: 7.02.04

Page: 6 of 6

Yoon SC, et al. Trends in renal transplantation in patients with human immunodeficiency virus infection: an analysis of the United States renal data system. Transplantation 2011 Apr 27;91(8):864-8.

*Key Article

KEY WORDS

Kidney Transplant, Renal Transplant

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently no National Coverage Determination (NCD) or Local Coverage Determination (LCD) for Kidney Transplantation. Please refer to the following NCD Complete Guide to Medicare Benefits Policy Manual for Transplantation - Inpatient Hospital Services at the following website for Medicare Members:

<https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c11.pdf>