

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	Bone Densitometry/Bone Density Studies
Policy Number	6.01.05
Category	Technology Assessment
Original Effective Date	11/19/99
Committee Approval Date	12/20/00, 04/19/01, 06/20/02, 10/16/02, 10/15/03, 11/18/04, 09/15/05, 02/06/06, 07/20/06, 05/17/07, 06/19/08, 09/17/09, 09/16/10, 09/15/11, 09/20/12, 09/19/13, 09/18/14, 09/17/15, 09/15/16, 09/21/17, 09/20/18, 09/19/19, 09/17/20, 09/16/21, 08/18/22, 08/17/23
Current Effective Date	08/17/23
Archived Date	N/A
Archive Review Date	N/A
Product Disclaimer	<ul style="list-style-type: none"> • <i>If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</i> • <i>If a commercial product (including an Essential Plan or Child Health Plus product), medical policy criteria apply to the benefit.</i> • <i>If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit.</i> • <i>If a Medicare product (including Medicare HMO-Dual Special Needs Program (DSNP) product) covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</i> • <i>If a Medicare HMO-Dual Special Needs Program (DSNP) product DOES NOT cover a specific service, please refer to the Medicaid Product coverage line.</i>

POLICY STATEMENT

- I. Based upon our criteria and assessment of the peer-reviewed literature, and in accordance with the New York Insurance Law, bone density measurements and tests are covered in accordance with the criteria of the Federal Medicare program and the National Institutes of Health (NIH). Bone mineral density (BMD) testing using dual-energy x-ray absorptiometry (DEXA or DXA), quantitative computed tomography (QCT), single-energy x-ray absorptiometry (SEXA), bone density of the heel using ultrasound, or quantitative ultrasound (QUS) is considered **medically appropriate** for the following patients, when the results will influence treatment decisions:
- All women aged 65 years and older, regardless of additional risk factors;
 - All men aged 70 years and older, regardless of additional risk factors;
 - Individuals previously diagnosed as having osteoporosis;
 - All post-menopausal women under age 65 years, who have one or more additional risk factors for osteoporotic fracture (please refer to description section for risk factors for osteoporosis in women) or who are at increased risk of osteoporosis, as determined by a formal clinical risk assessment tool (e.g., FRAX);
 - Men aged 50 to 70 years with clinical risk factors for osteoporotic fracture (please refer to description section for risk factors for osteoporosis in men);
 - Individuals on a prescribed drug regimen (e.g., anticonvulsants, aromatase inhibitors, cytotoxic drugs, Depo-Provera contraceptive injection, or hormone replacement therapy) posing a significant risk of osteoporosis;
 - Individuals receiving (or expecting to receive) long-term glucocorticoid therapy (e.g., glucocorticoids in a daily

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES

Policy Number: 6.01.05

Page: 2 of 12

- dose greater than or equal to 5 mg prednisone or equivalent for greater than or equal to three months);
- H. Individuals being monitored to assess the response to, or efficacy of an osteoporosis drug therapy approved by the U.S. Food and Drug Administration;
- I. Men and post-menopausal women aged 50 years and older with a current or prior hip or vertebral fracture;
- J. Men and post-menopausal women aged 50 years and older with other prior fractures and low bone mass (T-score between -1.0 and -2.5 at the femoral neck, total hip, or spine);
- K. Individuals with vertebral abnormalities, as demonstrated by an x-ray to be indicative of osteoporosis, osteopenia (low-bone mass), or vertebral fracture;
- L. Individuals with primary hyperparathyroidism;
- M. Individuals with suspected cases of secondary osteoporosis due to a broad range of disease states (e.g., hyperthyroidism, rheumatoid arthritis, or type 1 diabetes mellitus);
- N. Estrogen-deficient women at clinical risk for osteoporosis (please refer to description section for risk factors for osteoporosis in women);
- O. Post-menopausal women who have discontinued hormone replacement therapy within the past five years; and
- P. Men diagnosed with prostate cancer who have received either surgical or medical androgen deprivation therapy.

II. Follow-up testing:

The growing prevalence of osteoporosis, the management of the associated morbidity of osteoporotic fractures, and the monitoring of fracture prevention strategies present unique challenges. Unfortunately, there is little scientific data and, thus, a lack of consensus in peer-reviewed literature regarding the value, role, and interval of follow-up BMD testing. Changes in BMD may not be detected in less than two years of treatment because of the measurement technique, but a follow-up scan may be appropriate sooner in selected patients. Follow-up scans can help to detect treatment failure and secondary disease.

Follow-up BMD testing is indicated:

- A. Every two years (if at least 23 months have passed since the month the last BMD testing was performed), except for patients starting on Bisphosphonate therapy, for whom testing every three years after initiation of therapy is recommended; or
 - B. More frequently than every two (2) years, if medically necessary. Examples include, but are not limited to the following:
 - 1. The testing is performed to monitor individuals on long-term glucocorticoid (steroid) therapy of more than three months; or
 - 2. The testing is performed to obtain a confirmatory baseline (either central or peripheral) using a different BMD testing technique than was used in a patient's initial BMD test, to permit monitoring of the patient using the newer technique (e.g., if the initial test was performed using bone sonometry, and monitoring is anticipated using bone densitometry, a baseline measurement using bone densitometry is allowed); and
 - C. For women receiving estrogen replacement therapy (ERT), who should not be precluded from receiving follow-up or repeat BMD testing.
- III. Based upon our criteria and assessment of the peer-reviewed literature, screening for vertebral fracture with DEXA, SEXA or automated analysis of an existing computed tomography study is considered **investigational**.
- IV. Based upon our criteria and assessment of the peer-reviewed literature, pulse-echo ultrasound bone density measurement of the tibia and trabecular bone scoring for osteoporosis screening have not been medically proven to be effective and, therefore, are considered **investigational**.

Refer to Corporate Medical Policy #11.01.03 Experimental and Investigational Services

POLICY GUIDELINES

- I. Per the National Osteoporosis Foundation (NOF), BMD measurement is not routinely indicated in healthy men less than age 50 years or in pre-menopausal women.
- II. The decision to test for BMD should be based on an individual's risk profile. Testing is not indicated unless the

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES

Policy Number: 6.01.05

Page: 3 of 12

results would influence a treatment decision.

- III. Refer to the member's subscriber contract to confirm that the member's coverage is subject to the New York State BMD testing mandate.
- IV. Under the Affordable Care Act, non-grandfathered plans or policies are required to cover, in-network, without cost-sharing, the preventive services recommended by the United States Preventive Services Task Force (USPSTF), including osteoporosis screening for patients identified in Policy Statements I.A. and I.B. above.
- V. Whenever possible, use of central (axial skeleton) DEXA is preferred, as clinical evidence supports that this method has one of the lowest standard error rates in measurement and predictive accuracy. However, peripheral DEXA may be used as a substitute when technical problems preclude adequate imaging with a central DEXA machine or when a central DEXA machine is unavailable.
- VI. A DEXA study is representative of one or more sites; therefore, a study of multiple sites should be reported and will be processed as a single unit of service.
- VII. This policy addresses coverage for adults only. Requests for adolescents and children will be reviewed based on diseases and risk factors that are present. Per the National Osteoporosis Foundation, BMD measurement is not recommended in children or adolescents.

DESCRIPTION

The New York Insurance Law mandates that coverage be provided for bone density tests, as well as prescription drugs and devices that are approved by the FDA, for the detection and treatment of osteoporosis. The law provides that individuals qualifying for coverage shall, at a minimum, include individuals who have **ANY** of the following:

- I. A previous diagnosis or family history of osteoporosis;
- II. Symptoms or a condition indicative of the presence or significant risk of osteoporosis;
- III. A prescribed drug regimen posing a significant risk of osteoporosis;
- IV. Lifestyle factors posing a significant risk of osteoporosis; or
- V. Age, gender, and/or physiological characteristics posing a significant risk of osteoporosis.

Individuals are eligible for bone density measurements and tests in accordance with the criteria of the Federal Medicare program and the National Institutes of Health (NIH) for the detection of osteoporosis.

Defining osteoporosis by BMD:

The World Health Organization has established the following definitions, based on BMD of the spine, hip or forearm by DEXA. T scores are reported as standard deviations (SD):

- I. Normal: T-score at -1.0 and above (within 1 SD of a young healthy adult);
- II. Osteopenia (low bone mass): T-score between -1 to -2.5 (1 to 2.5 SD below that of a young healthy adult);
- III. Osteoporosis: T-score at or below -2.5 (2.5 SD or more below that of a young healthy adult); and
- IV. Severe osteoporosis: T-score of -2.5 or less with fragility fractures.

Although these definitions are necessary to establish the prevalence of osteoporosis, they should not be used as the sole determinant of treatment decisions.

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES

Policy Number: 6.01.05

Page: 4 of 12

Risk factors for osteoporosis in women:

There are certain symptoms, conditions, physiologic characteristics, and lifestyle factors indicative of a significant risk of osteoporosis. Per the NIH, risk factors for post-menopausal women under age 65 include, but are not limited to:

- I. Personal history of a non-traumatic fracture as an adult;
- II. Family history of osteoporosis (e.g., history of early, non-traumatic fracture in first-degree relative);
- III. Ethnicity - Caucasian and Asian women are at highest risk. African American and Latino women have a lower, but significant, risk;
- IV. Poor health/frailty;
- V. Cigarette smoking;
- VI. Body size – small, thin-boned women are at greater risk;
- VII. Extended estrogen deficiency;
- VIII. Age - Risk increases with age, as bones become less dense and weaker with age;
- IX. Eating disorders such as anorexia nervosa;
- X. Excessive use of alcohol;
- XI. Low calcium and vitamin D intake (lifelong);
- XII. Long-term use of certain medications, such as glucocorticoids;
- XIII. Sedentary lifestyle or extended bed rest; or
- XIV. Osteopenia.

The American College of Obstetricians and Gynecologists defines post-menopausal as the time in a woman's life when she stops having menstrual periods - specifically, when she has gone 12 consecutive months with no menstrual period. Menopause marks the end of the reproductive years that began in puberty.

Risk factors for osteoporosis in men:

There are certain symptoms, conditions, physiologic characteristics or lifestyle factors indicative of a significant risk of osteoporosis. Per the NIH, risk factors for osteoporosis in men include:

- I. Chronic diseases that affect the kidneys, lungs, stomach, and/or intestines or alters hormone levels;
- II. Long-term use of certain medications, such as glucocorticoids;
- III. Low levels of testosterone;
- IV. Unhealthy lifestyle habits (e.g., smoking, excessive alcohol use, low calcium intake, and inadequate physical exercise);
- V. Age – the older the individual, the greater the risk;
- VI. Race – Caucasian men appear to be at particularly high risk, but all men can develop osteoporosis.

RATIONALE

Several DEXA central bone densitometers have been approved by the FDA, such as the Norland XR 46 DXA (Central Bone Densitometer). Several bone ultrasonometers have been cleared for marketing by the FDA, such as Myriad's Soundscan (approved May 1998) and Hologic's Sahara Clinical Bone Sonometer (approved March 1998). To perform vertebral fracture assessment on DEXA devices, additional software is needed and must also have Section 510(k) marketing clearance from the FDA.

BMD studies:

While there are a number of tests available to assess bone mineral density, clinical evidence supports that DEXA of the hip or lumbar sacral spine and QCT have the lowest standard error in measurement and predictive accuracy. In addition, studies that show reduced fracture with treatment have used results from hip/spine DEXA machines.

The value of universal BMD screening, especially in peri-menopausal women, has not been established.

The U.S. Preventive Services Task Force (USPSTF) updated its recommendations in June 2018. Currently, the USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in women aged 65 years and older and in post-menopausal women younger than age 65 years who are at increased risk of osteoporosis, as

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES

Policy Number: 6.01.05

Page: 5 of 12

determined by a formal clinical risk assessment tool. The USPSTF concluded that the current evidence is insufficient to assess the balance of benefits and harms of screening for osteoporosis in men.

The National Osteoporosis Foundation (NOF) 2014 “Clinicians Guide to Prevention and Treatment of Osteoporosis” includes recommendation for BMD testing for women aged 65 years and older and for men aged 70 years and older, regardless of clinical risk factors. The guidelines also recommend BMD testing in: (a) post-menopausal women in the menopausal transition and men aged 50 to 69 years with clinical risk factors for fracture; (b) adults who have a fracture after age 50; and (c) adults who have a condition (e.g., rheumatoid arthritis) or are taking a medication (e.g., glucocorticoids in a daily dose ≥ 5 mg prednisone or equivalent for \geq three months) associated with low bone mass or bone loss. BMD measurement is not recommended in children or adolescents and is not routinely indicated in healthy young men or pre-menopausal women, unless there is a significant fracture history, or there are specific risk factors for bone loss. Follow-up testing is recommended one to two years after initiating medical therapy for osteoporosis and every two years thereafter.

The American College of Physicians (ACP) 2017 clinical practice guideline recommends against bone density monitoring during the five-year pharmacologic treatment period for osteoporosis in women. (Grade: weak recommendation; low quality evidence). Data from several studies showed women treated with bisphosphonates and other pharmacologic treatment benefited from reduced fractures with treatment, even if BMD did not increase. There is no evidence from randomized, controlled trials regarding how often to monitor BMD during osteoporosis treatment. There is moderate-quality evidence suggesting that most women do not need regular monitoring.

The Endocrine Society Clinical Practice Guideline: Osteoporosis in Men (2012) recommends testing higher-risk men (men age greater than or equal to 70 years and men aged 50 to 69 years who have risk factors [e.g., low body weight, prior fracture as an adult, smoking, etc.]) using central dual-energy x-ray absorptiometry. The guideline suggests measuring forearm DXA (1/3 or 33% radius) when spine or hip BMD cannot be interpreted and for men with hyperparathyroidism or receiving androgen deprivation therapy for prostate cancer.

Vertebral Fracture Assessment using DEXA:

Vertebral fractures are highly prevalent in the elderly populations, and epidemiologic studies have found that these fractures are associated with an increased risk of future spine or hip fractures independent of BMD. Only 20-30% of vertebral fractures are recognized clinically; the rest are discovered incidentally on lateral spine radiographs. Lateral spine x-rays have not been recommended as a component of risk assessment for osteoporosis, because of the cost, the radiation exposure, and the fact that the x-ray would require a separate procedure in addition to the BMD study. However, lateral spine images for vertebral fracture assessment (VFA) can be obtained using DEXA at the same time that a patient is undergoing assessment of BMD. Although VFA may be rapid and convenient, the images obtained are not of sufficient quality to establish the presence or absence of vertebral fractures. Literature suggests that densitometry may not accurately diagnose vertebral fractures in the population of interest (women with osteopenia or normal BMD). There is a lack of evidence that the test is accurate in detecting fractures among subjects without osteoporosis. Available evidence is insufficient to assess what health outcomes would result from VFA using DEXA as a screening test for osteoporosis. Evidence supporting the use of vertebral assessment using DEXA is not strong enough to reach conclusions about its effect on health outcomes.

Vertebral Fracture Assessment using Computed Tomography:

Vertebral fracture evaluation can be performed using a pre-existing computed tomography dataset via assistive augmented intelligence. It is used for the detection of osteoporosis as evidenced by vertebral compression fractures that are below the threshold for human detection. For example, the data set from a CTA scan of the chest performed for the diagnosis of pulmonary embolism could be subsequently analyzed by an algorithm created for identification of additional findings such as subtle osteoporotic vertebral fractures. There is currently limited research demonstrating the effectiveness of vertebral fracture assessment using CT with assistive augmented intelligence.

Pulse-echo Ultrasound:

Bone Index Finland, Ltd. received Section 510(k) approval from the FDA in May 2016 for Bindex BI-100 and Bindex BI-2 in January 2017. Bindex is a handheld pulse-echo ultrasound device used to measure apparent cortical bone thickness at the proximal tibia, to aid the physician in the diagnosis of osteoporosis. Use of pulse-echo ultrasonometry is being

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES

Policy Number: 6.01.05

Page: 6 of 12

promoted as a cost-effective screening option to determine appropriateness for DXA measurement. There is limited research, and no randomized, controlled trials, demonstrating the effectiveness of pulse-echo ultrasound as compared to DXA.

Trabecular Bone Scoring:

The trabecular bone score (TBS) is a diagnostic software tool that analyzes the internal structure of the lumbar spine bones. It is used in conjunction with dual-energy x-ray absorptiometry (DXA), which is an enhanced form of x-ray technology that is performed to measure bone mineral density (BMD). TBS uses standard DXA lumbar spine images to measure the structure of trabecular (cancellous) bone, providing additional information to be used in conjunction with BMD results in calculating fracture risk. Using a software program installed on a standard DXA computer, TBS provides an index of bone structure and quality without requiring additional radiation exposure or scan time. TBS received US FDA approval in 2012 for use as a complement to DXA analysis for the assessment of fracture risk and provides additional utility in fracture risk assessment among people with secondary causes of bone loss and fractures, such as type 2 diabetes. However, there is a lack of well-designed clinical trials including diverse populations in the peer-reviewed scientific literature addressing the impact of TBS on patient-specific health outcomes. Future prospective trials evaluating the use trabecular bone score in place of or in addition to established fracture prediction tools should report if long-term patient health outcomes are improved.

CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.
- Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN).

CPT Codes

Code	Description
76977	Ultrasound bone density measurement and interpretation, peripheral site(s), any method
77078	Computed tomography, bone mineral density study, 1 or more sites, axial skeleton (e.g., hips, pelvis, spine)
77080	Dual-energy X-ray absorptiometry (DXA), bone density study, 1 or more sites; axial skeleton (e.g., hips, pelvis, spine)
77081	Dual-energy X-ray absorptiometry (DXA), bone density study, 1 or more sites; appendicular skeleton (peripheral) (e.g., radius, wrist, heel)
77085 (E/I)	Dual-energy X-ray absorptiometry (DXA), bone density study, 1 or more sites; axial skeleton (e.g., hips, pelvis, spine), including vertebral fracture assessment
77086 (E/I)	Vertebral fracture assessment via dual-energy X-ray absorptiometry (DXA)
77089 (E/I)	Trabecular bone score(TBS), structural condition of the bone microarchitecture; using dual X-ray absorptiometry (DXA) or other imaging data on gray-scale variogram, calculation, with interpretation and report on fracture-risk
77090 (E/I)	Trabecular bone score (TBS), structural condition of the bone microarchitecture; technical preparation and transmission of data for analysis to be performed elsewhere

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES**Policy Number: 6.01.05****Page: 7 of 12**

Code	Description
77091 (E/I)	Trabecular bone score (TBS), structural condition of the bone microarchitecture; technical calculation only
77092 (E/I)	Trabecular bone score (TBS), structural condition of the bone microarchitecture; interpretation and report on fracture-risk only by other qualified health care professional
0508T (E/I)	Pulse-echo ultrasound bone density measurement resulting in indicator of axial bone mineral density, tibia
0554T (E/I)	Bone strength and fracture risk using finite element analysis of functional data, and bone-mineral density, utilizing data from a computed tomography scan; retrieval and transmission of the scan data, assessment of bone strength and fracture risk and bone mineral density, interpretation and report
0555T (E/I)	retrieval and transmission of the scan data
0556T (E/I)	assessment of bone strength and fracture risk and bone mineral density
0557T (E/I)	interpretation and report
0691T (E/I)	Automated analysis of an existing computed tomography study for vertebral fracture(s), including assessment of bone density when performed, data preparation, interpretation, and report (Do not report 0691T in conjunction with 71250, 71260, 71270, 71271, 71275, 72125, 72126, 72127, 72128, 72129, 72130, 72131, 72132, 72133, 72191, 72192, 72193, 72194, 74150, 74160, 74170, 74174, 74175, 74176, 74177, 74178, 74261, 74262, 74263, 75571, 75572, 75573, 75574, 75635, 78814, 78815, 78816, 0554T, 0555T, 0556T, 0557T, 0558T)
0743T (E/I)	Bone strength and fracture risk using finite element analysis of functional data and bone-mineral density (BMD), with concurrent vertebral fracture assessment, utilizing data from a computed tomography scan, retrieval and transmission of the scan data, measurement of bone strength and BMD and classification of any vertebral fractures, with overall fracture risk assessment, interpretation and report (<i>Effective 01/01/2023</i>)

Copyright © 2023 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
G0130	Single energy x-ray absorptiometry (SEXA) bone density study, one or more sites; appendicular skeleton (peripheral) (e.g., radius, wrist, heel)

ICD10 Codes

Code	Description
E21.0-E21.5	Hyperparathyroidism and other disorders of parathyroid gland (code range)
E24.0-E24.9	Cushing's syndrome (code range)
E28.310- E28.319	Premature menopause (code range)
E28.39	Other primary ovarian failure
E29.1	Testicular hypofunction

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES**Policy Number: 6.01.05****Page: 8 of 12**

Code	Description
E34.2	Ectopic hormone secretion, not elsewhere classified
E89.40	Asymptomatic postprocedural ovarian failure
E89.41	Symptomatic postprocedural ovarian failure
E89.5	Postprocedural testicular hypofunction
N91.0-N91.2	Amenorrhea (code range)
N92.4	Excessive bleeding in the premenopausal period
N95.0-N95.9	Menopausal and other perimenopausal disorders (code range)
M48.50XA- M48.58XA	Collapsed vertebra, not elsewhere classified (code range)
M80.00XA	Age-related osteoporosis with current pathological fracture, unspecified site, initial encounter for fracture
M80.011A	Age-related osteoporosis with current pathological fracture, right shoulder, initial encounter for fracture
M80.012A	Age-related osteoporosis with current pathological fracture, left shoulder, initial encounter for fracture
M80.019A	Age-related osteoporosis with current pathological fracture, unspecified shoulder, initial encounter for fracture
M80.031A	Age-related osteoporosis with current pathological fracture, right forearm, initial encounter for fracture
M80.032A	Age-related osteoporosis with current pathological fracture, left forearm, initial encounter for fracture
M80.039A	Age-related osteoporosis with current pathological fracture, unspecified forearm, initial encounter for fracture
M80.041A	Age-related osteoporosis with current pathological fracture, right hand, initial encounter for fracture
M80.042A	Age-related osteoporosis with current pathological fracture, left hand, initial encounter for fracture
M80.049A	Age-related osteoporosis with current pathological fracture, unspecified hand, initial encounter for fracture
M80.051A	Age-related osteoporosis with current pathological fracture, right femur, initial encounter for fracture
M80.052A	Age-related osteoporosis with current pathological fracture, left femur, initial encounter for fracture
M80.059A	Age-related osteoporosis with current pathological fracture, unspecified femur, initial encounter for fracture
M80.061A	Age-related osteoporosis with current pathological fracture, right lower leg, initial encounter for fracture

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES**Policy Number: 6.01.05****Page: 9 of 12**

Code	Description
M80.062A	Age-related osteoporosis with current pathological fracture, left lower leg, initial encounter for fracture
M80.069A	Age-related osteoporosis with current pathological fracture, unspecified lower leg, initial encounter for fracture
M80.071A	Age-related osteoporosis with current pathological fracture, right ankle and foot, initial encounter for fracture
M80.072A	Age-related osteoporosis with current pathological fracture, left ankle and foot, initial encounter for fracture
M80.079A	Age-related osteoporosis with current pathological fracture, unspecified ankle and foot, initial encounter for fracture
M80.08XA	Age-related osteoporosis with current pathological fracture, vertebra(e), initial encounter for fracture
M80.811A	Other osteoporosis with current pathological fracture, right shoulder, initial encounter for fracture
M80.812A	Other osteoporosis with current pathological fracture, left shoulder, initial encounter for fracture
M80.819A	Other osteoporosis with current pathological fracture, unspecified shoulder, initial encounter for fracture
M80.831A	Other osteoporosis with current pathological fracture, right forearm, initial encounter for fracture
M80.832A	Other osteoporosis with current pathological fracture, left forearm, initial encounter for fracture
M80.839A	Other osteoporosis with current pathological fracture, unspecified forearm, initial encounter for fracture
M80.841A	Other osteoporosis with current pathological fracture, right hand, initial encounter for fracture
M80.842A	Other osteoporosis with current pathological fracture, left hand, initial encounter for fracture
M80.849A	Other osteoporosis with current pathological fracture, unspecified hand, initial encounter for fracture
M80.851A	Other osteoporosis with current pathological fracture, right femur, initial encounter for fracture
M80.852A	Other osteoporosis with current pathological fracture, left femur, initial encounter for fracture
M80.859A	Other osteoporosis with current pathological fracture, unspecified femur, initial encounter for fracture
M80.861A	Other osteoporosis with current pathological fracture, right lower leg, initial encounter for fracture

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES**Policy Number: 6.01.05****Page: 10 of 12**

Code	Description
M80.862A	Other osteoporosis with current pathological fracture, left lower leg, initial encounter for fracture
M80.869A	Other osteoporosis with current pathological fracture, unspecified lower leg, initial encounter for fracture
M80.871A	Other osteoporosis with current pathological fracture, right ankle and foot, initial encounter for fracture
M80.872A	Other osteoporosis with current pathological fracture, left ankle and foot, initial encounter for fracture
M80.879A	Other osteoporosis with current pathological fracture, unspecified ankle and foot, initial encounter for fracture
M80.88XA	Other osteoporosis with current pathological fracture, vertebra(e), initial encounter for fracture
M81.0-M81.8	Osteoporosis without current pathological fracture (code range)
M84.48XA	Pathological fracture, other site, initial encounter for fracture
M84.58XA	Pathological fracture in neoplastic disease, other specified site, initial encounter for fracture
M84.68XA	Pathological fracture in other disease, other site, initial encounter for fracture
M85.80	Other specified disorders of bone density and structure, unspecified site
M85.811- M85.819	Other specified disorders of bone density and structure, shoulder (code range)
M85.821- M85.829	Other specified disorders of bone density and structure, upper arm (code range)
M85.831- M85.839	Other specified disorders of bone density and structure, forearm (code range)
M85.841- M85.849	Other specified disorders of bone density and structure, hand (code range)
M85.851- M85.859	Other specified disorders of bone density and structure, thigh (code range)
M85.861- M85.869	Other specified disorders of bone density and structure, lower leg (code range)
M85.871- M85.879	Other specified disorders of bone density and structure, ankle and foot (code range)
M85.88	Other specified disorders of bone density and structure, other site
M85.89	Other specified disorders of bone density and structure, multiple sites
Q96.0-Q96.9	Turner's syndrome (code range)
R29.890	Loss of height
Z08	Encounter for follow-up examination after completed treatment for malignant neoplasm

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES

Policy Number: 6.01.05

Page: 11 of 12

Code	Description
Z09	Encounter for follow-up examination after completed treatment for conditions other than malignant neoplasm
Z13.820	Encounter for screening for osteoporosis
Z78.0	Asymptomatic menopausal state
Z79.51-Z79.52	Long term (current) use of steroids (code range)
Z87.310	Personal history of (healed) osteoporosis fracture
Z87.311	Personal history of (healed) other pathological fracture
Z87.312	Personal history of (healed) stress fracture
Z90.721- Z90.722	Acquired absence of ovaries (code range)
Z90.79	Acquired absence of other genital organ(s)

REFERENCES

Adler RA. Update on osteoporosis in men. Best Pract Res Clin Endocrinol Metab 2018 Oct;32(5):759-772.

*Agency for Healthcare Research and Quality. Osteoporosis in postmenopausal women: diagnosis and monitoring. Evidence Report/Technology Assessment 2001 Apr 9;28.

*Agency for Healthcare Research and Quality (AHRQ). Guide to Clinical Preventive Services. Recommendations of the U.S. Preventive Services Task Force. Screening for osteoporosis. 2021 November.

[<https://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/osteoporosis-screening>] accessed 07/24/23.

*American College of Obstetricians and Gynecologists. ACOG guidelines for the clinical management of osteoporosis. 2004 Mar 15 [<http://www.aafp.org/afp/2004/0315/p1558.html>] accessed 07/24/23.

*American College of Obstetricians and Gynecologists. Osteoporosis. ACOG Practice Bulletin No. 50, Obstet Gynecol 2004 Jan;103(1):203-16.

Diem SJ, et al. Screening for osteoporosis in older men: operating characteristics of proposed strategies for selecting men for BMD testing. J Gen Intern Med 2017 Nov;32(11):1235-1241.

El-Hajj Fuleihan G, et al. Worldwide fracture prediction. J Clin Densitom 2017 Jul 19;20(3): 397-424.

Ginther JP, et al. Adding VFA to DXA identifies fracture risk in a way not duplicated by other measures. Endocr Pract 2017 Dec;23(12):1375-1378.

Hans D, et al. Long-term effect of denosumab on bone microarchitecture as assessed by tissue thickness-adjusted trabecular bone score in postmenopausal women with osteoporosis: results from FREEDOM and its open-label extension. Osteoporos Int 2023 Jun;34(6):1075-1084.

Kim DK, et al. Effect of androgen-deprivation therapy on bone mineral density in patients with prostate cancer: a systematic review and meta-analysis. J Clin Med 2019 Jan 18;8(1):E113.

Löffler MT, et al. Automatic opportunistic osteoporosis screening in routine CT: improved prediction of patients with prevalent vertebral fractures compared to DXA. European Radiology 2021;31:6069–6077.

Merlijn T, et al. The effect of a screening and treatment program for the prevention of fractures in older women: a randomized pragmatic trial. Journal of Bone and Mineral Research 2019 Nov;34(11):1993–2000.

Medical Policy: BONE DENSITOMETRY/ BONE DENSITY STUDIES

Policy Number: 6.01.05

Page: 12 of 12

National Institutes of Health Osteoporosis and Related Bone Diseases National Resource Center. Osteoporosis overview. 2019 Oct [http://www.niams.nih.gov/Health_Info/Bone/Osteoporosis/overview.asp] accessed 07/24/23.

National Institute of Health Osteoporosis and Related Bone Diseases National Resource Center. Osteoporosis in men. 2018 Oct. [http://www.niams.nih.gov/Health_Info/Bone/Osteoporosis/men.asp] accessed 07/24/23.

National Osteoporosis Foundation (NOF) / International Society for Clinical Densitometry (ISCD) FRAX Implementation Guide [<https://iscd.org/learn/resources/fracture-risk-models/>] accessed 07/24/23.

O'Donnell S, et al. Screening, prevention and management of osteoporosis among Canadian adults. Health Promot Chronic Dis Prev Can 2018 Dec;38(12):445-454.

Pan Y, et al. Automatic opportunistic osteoporosis screening using low-dose chest computed tomography scans obtained for lung cancer screening. European Radiology 2020;30:4107-4116.

Pouilles J-M, et al. TBS in early postmenopausal women with severe vertebral osteoporosis. Bone 2021;142:115698.

*Qaseem A, et al. Treatment of low bone density or osteoporosis to prevent fracture in men and women : a clinical practice guideline update from the American College of Physicians. Ann Intern Med 2017 Jun 6;166(11):818-839.

Salari N, et al. Global prevalence of osteoporosis among the world older adults: a comprehensive systematic review and meta-analysis. Journal of Orthopaedic Surgery and Research 2021;16:669.

*State of New York Insurance Law §7657, Amendment 11723, effective 2003 Jan 1.

US Preventive Services Task Force Screening for osteoporosis: U.S. Preventive Services Task Force recommendation statement. JAMA 2018 Jun 26;319(24):2521-2531.

Ward RJ, et al. Expert Panel on musculoskeletal imaging: ACR appropriateness criteria osteoporosis and bone mineral density. J Am Coll Radiol 2017 May;14(5S):S189-S202.

White Van Gompel EC, et al. Incidence and predictors of repeat bone mineral densitometry: a longitudinal cohort study. J Gen Intern Med 2017 Oct;32(10):1090.

World Health Organization Fracture Risk Assessment Tool [<https://www.sheffield.ac.uk/FRAX/>] accessed 07/24/23.

Yang J. et al. Identification of prevalent vertebral fractures using Vertebral Fracture Assessment (VFA) in asymptomatic postmenopausal women: A systematic review and meta-analysis. Bone 2020;136:doi115358.

*Key Article

KEY WORDS

BMD, bone mineral density, DEXA, Fracture Risk Assessment Tool, FRAX, vertebral fracture assessment, morphometric x-ray absorptiometry.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a National Coverage Determination (NCD) for Bone (Mineral) Density Studies (150.3). Please refer to the following NCD website for Medicare Members: [<https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=256&ncdver=2&bc=AgAAgAAAAAAA&>]