

# MEDICAL POLICY

Medical Policy Title	Evoked Potentials
Policy Number	2.01.27
Current Effective Date	May 21, 2026
Next Review Date	May 2027

Our medical policies are guides to evaluate technologies or services for medical necessity. Criteria are established through the assessment of evidence based, peer-reviewed scientific literature, and national professional guidelines. Federal and state law(s), regulatory mandates and the member's subscriber contract language are considered first in the determination of a covered service. ([Link to Product Disclaimer](#))

## POLICY STATEMENT(S)

- I. Visual-evoked potentials are considered **medically appropriate** for **ANY** of the following indications:
  - A. To diagnose and monitor multiple sclerosis;
  - B. To localize the cause of a visual field defect that is not explained by lesions seen on CT, or MRI, metabolic disorder, or infectious disease;
  - C. To diagnose or evaluate deficits or damage to the visual system of infants or unresponsive/nonverbal patients.
- II. Somatosensory-evoked potentials are considered **medically appropriate** for **ANY** of the following indications:
  - A. To assess any decline that may be considered emergent to surgery in unconscious spinal cord injury patients who show specific structural damage to the somatosensory system and who are candidates for emergency spinal cord surgery;
  - B. To diagnose and monitor multiple sclerosis;
  - C. To evaluate patients with suspected brain death;
  - D. To diagnose unexplained myelopathy;
  - E. To localize the cause of a neurologic deficit seen on exam and not explained by lesions on CT or MRI.
- III. Auditory-evoked potentials are considered **medically appropriate** for **ANY** of the following indications:
  - A. To evaluate brainstem function in acquired metabolic disorders;
  - B. To assess recovery of brainstem function after a lesion compressing the brainstem has been surgically removed;
  - C. To localize the cause of a neurologic deficit seen on exam, not explained by lesions seen on computed tomography (CT) scan or magnetic resonance imaging (MRI);

## Medical Policy: Evoked Potentials

Policy Number: 02.01.27

Page: 2 of 13

- D. To diagnose and monitor demyelinating and degenerative diseases affecting the brain stem (e.g., multiple sclerosis, central pontine myelinolysis, olivopontocerebellar degeneration, and others);
  - E. To diagnose lesions in the auditory system;
  - F. To evaluate the irreversibility of coma or brain death, along with an electroencephalogram (EEG);
  - G. For children under age five (5), to determine the type and degree of hearing problems or to determine the developed status of nerves.
- IV. Intra-operative neurophysiologic monitoring is **medically appropriate** during high-risk thyroid or parathyroid surgery, or during spinal, intracranial, or vascular procedures.
- V. All other indications for intra-operative neurophysiologic monitoring are considered **not medically necessary**.
- VI. Vestibular evoked myogenic potentials (cVEMPs and oVEMPs) are considered **investigational**.
- VII. Intra-operative monitoring of visual-evoked potentials is considered **investigational**.
- VIII. Visual-evoked potential testing for the diagnosis and evaluation of glaucoma is considered **investigational**.
- IX. The use of transcranial magnetic stimulation to elicit motor-evoked potentials is considered **investigational**.

### RELATED POLICIES

Corporate Medical Policy

11.01.03 Experimental or Investigational Services

### POLICY GUIDELINE(S)

Not Applicable

### DESCRIPTION

Evoked potentials (EP) are responses (electrical signals) produced by the nervous system in response to a stimulus. These computerized tests help diagnose nerve disorders, locate the site of nerve damage, and evaluate the individual's condition after treatment or during surgery. There are several types of EP tests. Each uses mild stimulus to cause the nerves to react and send a message to the brain. Electrodes placed on the skin surface record how the brain and spinal cord respond to stimulus. The responses are analyzed by a computer and printed as a waveform pattern. The wave pattern may reveal certain problems and show where any damage is located along the nerve pathway(s) being tested. EPs can be further broken down into the following categories, according to the type of stimulation used:

Somatosensory-Evoked Potentials

## **Medical Policy: Evoked Potentials**

**Policy Number: 02.01.27**

**Page: 3 of 13**

Somatosensory-evoked potentials (SSEPs) are electrical waves that are generated by the response of sensory neurons to stimulation. Peripheral nerves, such as the median, ulnar or tibial nerves, are typically stimulated, but in some situations the spinal cord may be stimulated directly. Recording is done either cortically or at the level of the spinal cord above the surgical procedure.

### Auditory-Evoked Potentials

Auditory-evoked potentials, also called auditory brainstem response (ABR), are electrophysiologic measures of auditory function that utilizes responses produced by the auditory nerve and brainstem and helps differentiate sensory from neural hearing loss. The response is the waveform averaged over many auditory clicks.

Brainstem auditory-evoked potentials (BAEPs) are generated in response to auditory clicks and can define the functional status of the auditory nerve. Surgical resection of a cerebellopontine angle tumor, such as an acoustic neuroma, places the auditory nerves at risk, and BAEPs have been extensively used to monitor auditory function during these procedures. The clinical utility of BAEP over standard auditory testing is due to BAEP's characteristics: (1) BAEP's resistance to alteration by systemic metabolic abnormalities, medications or pronounced changes in the state of consciousness of the patient; and (2) the close association of BAEP waveform abnormalities to underlying structural pathology. BAEP has been proven effective for differentiating conductive from sensory hearing loss, for detecting tumors and other disease states affecting central auditory pathways (e.g., acoustic neuromas, subclinical lesions in MS), and for non-invasively detecting hearing loss in patients who cannot cooperate with subjective auditory testing (e.g., infants, comatose patients). BAEP is the test of choice to assess hearing in infants and young children. It is most useful for following asphyxia, hyperbilirubinemia, intracranial hemorrhage, or meningoencephalitis or for assessing an infant who has trisomy. BAEP also is useful in the assessment of MS or other demyelinating condition, coma, or hysteria. Audiometric analysis using multiple sound frequencies is usually preferred over BAEP for testing hearing in cooperative patients who are able to report when sounds are heard.

### Vestibular Evoked Myogenic Potentials

Vestibular-evoked myogenic potentials (VEMPs) testing is described as an alternative method of assessing otolith organs and their function, with a goal of diagnosing suspected vestibular disorders. VEMPs are performed by repeatedly stimulating the ears using loud sounds such as clicks and tone bursts or bone vibration. These sounds/vibrations create a pressure stimulus. VEMPs are recorded using slightly modified standard evoked potential equipment. The cervical VEMP (cVEMP) pathway begins in the saccule, that travels through the inferior vestibular nerve, vestibular nucleus, medial vestibulospinal tract, the accessory nucleus, the eleventh nerve, and then to a contracted ipsilateral sternocleidomastoid muscle where it is recorded via surface electrode. An abnormality in any part of the pathway is thought to affect the response.

Ocular VEMPs (oVEMPs) stimulate the otolithic membrane, traveling to the vestibule to trigger the utricle and the vestibular ocular reflex causing contraction of the extraocular muscle and is measured by surface electrodes placed under the contralateral eye during an upward gaze. VEMPs have been investigated for the diagnosis of vestibular neuritis, benign paroxysmal positional vertigo, vestibular

## Medical Policy: Evoked Potentials

Policy Number: 02.01.27

Page: 4 of 13

schwannoma, Meniere disease, vestibular migraine, superior semicircular canal dehiscence (SCDS), and other vestibular disorders.

### Visual-Evoked Potentials

Visual-evoked potentials (VEPs) are used to track visual signals from the retina to the occipital cortex. VEP monitoring has been used for surgery on lesions near optic chiasm. However, intra-operatively recorded VEPs are very difficult to interpret, due to their sensitivity to anesthesia, temperature, and blood pressure.

### Motor-Evoked Potentials

Motor-evoked potentials (MEPs) are elicited by either electrical or magnetic stimulation of the motor cortex or the spinal cord. Transcranial electrical stimulation involves stimulation of the motor cortex via electrodes placed on the scalp, or, if the brain is exposed by a craniotomy, placed directly on the brain surface. Magnetic stimulation delivers a pulsed magnetic field over the scalp in the region of the primary motor cortex. Magnetic stimulation is regarded as unsuitable for intra-operative monitoring because it is more sensitive to anesthesia.

### Intra-Operative Neurophysiologic Monitoring

Intra-operative neurophysiologic monitoring (IONM) describes a variety of procedures used to monitor the integrity of neural pathways during high-risk neurosurgical, orthopedic, and vascular surgeries. It involves the detection of electrical signals produced by the nervous system in response to sensory or electrical stimuli, to provide information about the functional integrity of neuronal structures. Different methodologies include, but may not be limited to, SSEPs, MEPs using transcranial electrical stimulation, BAEPs, electromyography (EMG) of cranial nerves, electroencephalography, and electrocorticography. IONM has been utilized in attempts to minimize neurological morbidity from operative manipulations. The goal of such monitoring is to identify changes in brain, spinal cord, and peripheral nerve function prior to irreversible damage. Intraoperative monitoring also has been effective in localizing anatomical structures, including peripheral nerves and sensorimotor cortex, which helps guide the surgeon during dissection. SSEP has been the standard of intraoperative monitoring, with excellent ability to assess dorsal column and lateral sensory tract function; it may also be able to detect changes in function of anterior motor tracts by stimulating mixed sensorimotor peripheral nerves. However, significant motor deficits have been seen in individuals undergoing spinal surgery, despite normal SSEPs. MEPs were developed to better monitor the motor neurophysiological pathways.

In individuals with pre-operative spinal cord compromise, MEPs may be present when SSEPs are absent or ill-defined. This is because MEPs and SSEPs are conducted in different spinal cord pathways and have different blood supplies. Consequently, being able to perform MEP monitoring makes spinal cord monitoring possible in cases where SSEP signals are unobtainable. In the operating room, transcranial electrical stimulation is preferable to transcranial magnetic stimulation, because the electrical stimulus is more reproducible.

### **SUPPORTIVE LITERATURE**

## **Medical Policy: Evoked Potentials**

**Policy Number: 02.01.27**

**Page: 5 of 13**

There is sufficient data published in the medical literature to conclude that measurement of evoked potentials (EPs) and intra-operative monitoring of EPs, in appropriate situations, improves health outcome. Improved health outcomes have been achieved outside the investigational setting.

### Visual-Evoked Potentials

Clinical studies have demonstrated a statistically significant association between abnormal VEPs and an increased risk of developing clinically definite multiple sclerosis (CDMS). In these studies, individuals with suspected multiple sclerosis (MS) were 2.5 to 9 times as likely to develop CDMS as patients with normal VEPs. VEP sensitivities ranged from 25% to 83% (Gronseth 2000). VEPs improved the ability to predict which MS suspects will develop CDMS by as much as 29%.

Measurement of visual-evoked responses (VERs) is the primary means of objectively testing vision in infants and young children. VER measurements are useful in infants and young children suspected of having disorders of the visual system, where the child is too young to report differences in color vision or to undergo assessment of visual fields and visual acuity. Lesions affecting the visual pathways can be localized by noting the presence of decreased amplitudes or increased latencies of VERs, and by determining whether VER abnormalities involve one or both eyes.

Several small studies (Pillai 2013; Mousa 2014; Jha 2017; Waisbourd 2017) have investigated the use of VEP technology to differentiate between normal, healthy eyes and eyes with early-to-advanced visual field loss resulting from glaucoma. The authors indicated that VEP signals may discriminate between normal eyes and glaucomatous eyes. However, larger studies are needed to confirm these findings. Additionally, VEP has not been shown to be superior to standard visual field testing in the diagnosis of glaucoma or management of clinical outcomes.

Published studies mostly assess how cVEMP and oVEMP change with various disease states as opposed to assessing the diagnostic accuracy or determination of appropriate populations. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

### Intra-Operative Neurophysiologic Monitoring

For individuals who receive IONM while undergoing thyroid or parathyroid surgery due to high risk of injury to the recurrent laryngeal nerve (RLN), the evidence includes a large, randomized, controlled trial (RCT) and systematic reviews. Relevant outcomes are morbid events, functional outcomes, and quality of life. The strongest evidence on neurophysiologic monitoring derives from an RCT of 1,000 patients undergoing thyroid surgery (Barczynski 2009). This RCT found a significant reduction in RLN injury in patients at high risk for injury. "High risk" in this trial was defined as surgery for cancer, thyrotoxicosis, retrosternal or giant goiter, or thyroiditis. The high-risk category may also include patients with prior thyroid or parathyroid surgery or total thyroidectomy. A low volume of surgeries might also contribute to a higher risk for RLN injury. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Cozzi et al (2023) reported on a systematic review of 164 studies examining intraoperative neurophysiologic monitoring during thyroid surgery. The overall rates of temporary and permanent recurrent laryngeal nerve injury were 3.15% and 0.422%, respectively, across all procedures. In cases where intraoperative neurophysiologic monitoring was utilized, these rates were 3.29% and

## Medical Policy: Evoked Potentials

Policy Number: 02.01.27

Page: 6 of 13

0.409% for permanent injury. In contrast, the rates were 3.16% and 0.463%, respectively. The pooled rates of temporary recurrent laryngeal nerve injury were 2.48% for intermittent intraoperative neurophysiologic monitoring and 2.913% for continuous monitoring, while the pooled rates for definitive injury rates, the pooled rates were 0.395% and 0.40%, respectively. The authors noted that pooled rates had overlapping 95% confidence intervals (CI), concluding that intraoperative neurophysiologic monitoring does not significantly impact the rates of temporary or definitive recurrent laryngeal nerve injury following thyroidectomy.

### Vestibular Evoked Myogenic Potential Testing

A cohort study by Verrecchia et al (2019) evaluated oVEMP testing in individuals experiencing dizziness. The study included ten patients diagnosed with superior canal dehiscence syndrome and 135 individuals with dizziness that were not already diagnosed as superior canal dehiscence syndrome (SCDS). The study included four oVEMP parameters (amplitude, latency, amplitude asymmetry ratio, and interaural latency difference). oVEMP was able to identify all subjects affected by SCDS, in a large cohort of unselected dizzy patients. Of the parameters, the amplitude to air-conducted stimulation had the greatest diagnostic accuracy (area under the curve score, 0.96). A cutoff point of 16.7  $\mu\text{V}$  resulted in a sensitivity of 100% and specificity of 89%. The available evidence has indicated that the use of VEMP tests to evaluate suspected vestibular disorders is at a very early stage of development. Standardization of procedures and studies on the diagnostic accuracy of these procedures in the appropriate populations are needed.

### **PROFESSIONAL GUIDELINE(S)**

The 2017 Practice Guidelines from American Academy of Neurology (AAN) (reaffirmed in 2024) assessed the diagnostic value of vestibular evoked myogenic potential testing in individuals with vestibular symptoms. The conditions of interest included superior canal dehiscence syndrome, vestibular neuritis or migraine, Meniere disease, and benign paroxysmal positional vertigo (BPPV). The evidence for testing in BPPV was drawn from two class III studies, neither of which presented sufficient diagnostic value of vestibular evoked myogenic potential testing for the treatment to be recommended (level C evidence).

In 2020, the Scoliosis Research Society published an information statement on neurophysiologic monitoring during spinal deformity surgery. The Society concluded that neurophysiologic monitoring facilitates early detection of complications and may help prevent postoperative neurologic injury. It is regarded as optimal care when the spinal cord is at risk which warrants a strong recommendation unless contraindications are present period the standard method for intraoperative monitoring should include transcranial motor evoked potentials and somatosensory evoked potentials with or without additional electromyographic monitoring.

In 2022, the American Head and Neck Society Endocrine Surgery Section and the International Neural Monitoring Study Group published a clinical review on intraoperative nerve monitoring during pediatric thyroid surgery. The review indicated that intraoperative neurophysiological monitoring can be considered for all pediatric thyroid surgeries. Procedures for which monitoring may be particularly beneficial include total thyroidectomy, hemithyroidectomy in cases where the contralateral vocal cord is paralyzed, and reoperative surgeries.

## Medical Policy: Evoked Potentials

Policy Number: 02.01.27

Page: 7 of 13

In 2023, the American Association of Neuromuscular & Electrodiagnostic Medicine (AANEM) recommended that intraoperative sensory-evoked potentials have demonstrated usefulness for monitoring of spinal cord, brainstem, and brain sensory tracts. The AANEM stated that intraoperative somatosensory-evoked potential monitoring is indicated for select spine surgeries in which there is a risk of additional nerve root or spinal cord injury. Indications for somatosensory-evoked potential monitoring may include, but are not limited to, complex, extensive, or lengthy procedures.

### REGULATORY STATUS

Not Applicable

### CODE(S)

- Codes may not be covered under all circumstances.
- Code list may not be all inclusive (AMA and CMS code updates may occur more frequently than policy updates).
- (E/I)=Experimental/Investigational
- (NMN)=Not medically necessary/appropriate

### CPT Codes

Code	Description
92517 (E/I)	Vestibular evoked myogenic potential (VEMP) testing, with interpretation and report, cervical (cVEMP)
92518 (E/I)	Vestibular evoked myogenic potential (VEMP) testing, with interpretation and report, ocular (oVEMP)
92519 (E/I)	Vestibular evoked myogenic potential (VEMP) testing, with interpretation and report, cervical (cVEMP) and ocular (oVEMP)
92650	Auditory evoked potentials; screening of auditory potential with broadband stimuli, automated analysis
92651	Auditory evoked potentials; for hearing status determination, broadband stimuli, with interpretation and report
92652	Auditory evoked potentials; for threshold estimation at multiple frequencies, with interpretation and report
92653	Auditory evoked potentials; neurodiagnostic, with interpretation and report
95925-95927, 95938	Somatosensory-evoked potentials (code range)
95928-95929, 95939	Central motor evoked potential study (transcranial motor stimulation) (code range)
95930	Visual-evoked potential (VEP) checkerboard or flash testing, central nervous system except glaucoma, with interpretation and report

**Medical Policy: Evoked Potentials****Policy Number: 02.01.27****Page: 8 of 13**

Code	Description
95940	Continuous intraoperative neurophysiology monitoring in the operating room, one on one monitoring requiring personal attendance, each 15 minutes (List separately in addition to code for primary procedure)
95941	Continuous intraoperative neurophysiology monitoring, from outside the operating room (remote or nearby) or for monitoring of more than one case while in the operating room, per hour (List separately in addition to code for primary procedure)
0333T (E/I)	Visual evoked potential, screening of visual acuity, automated, with report
0464T (E/I)	Visual evoked potential, testing for glaucoma, with interpretation and report

Copyright © 2026 American Medical Association, Chicago, IL

**HCPCS Codes**

Code	Description
G0453	Continuous intraoperative neurophysiology monitoring, from outside the operating room (remote or nearby) per patient (attention directed exclusively to one patient) each 15 minutes (List separately in addition to code for primary procedure)

**ICD10 Codes**

Code	Description
C71.6	Malignant neoplasm of cerebellum
C79.31	Secondary malignant neoplasm of brain
D33.0-D33.2	Benign neoplasm of brain (code range)
D33.3	Benign neoplasm of cranial nerves
D43.0-D43.2	Neoplasm of uncertain behavior of brain (code range)
D43.4	Neoplasm of uncertain behavior of spinal cord
D49.6	Neoplasm of unspecified behavior of brain
H40.001- H40.9 (E/I)	Glaucoma (code range)
H53.411- H53.419	Scotoma involving central area (code range)
H53.421- H53.429	Scotoma of blind spot area (code range)
H53.431- H53.439	Sector or arcuate defects (code range)
H53.451- H53.459	Other localized visual field defect (code range)

**Medical Policy: Evoked Potentials****Policy Number: 02.01.27****Page: 9 of 13**

<b>Code</b>	<b>Description</b>
H53.461- H53.469	Homonymous bilateral field defects (code range)
H53.47	Heteronymous bilateral field defects
H53.481- H53.489	Generalized contraction of visual field (code range)
I63.031- I63.039	Cerebral infarction due to thrombosis of carotid artery (code range)
I63.131- I63.139	Cerebral infarction due to embolism of carotid artery (code range)
I63.231- I63.239	Cerebral infarction due to unspecified occlusion or stenosis of carotid arteries (code range)
I65.21-I65.29	Occlusion and stenosis of carotid artery (code range)
I71.00-I71.03	Dissection of aorta (code range)
M40.00- M40.05	Postural kyphosis (code range)
M40.202- M40.209	Unspecified kyphosis (code range)
M40.292- M40.299	Other kyphosis (code range)
M40.30- M40.37	Flatback syndrome (code range)
M40.40- M40.57	Lordosis (code range)
M41.00- M41.27	Idiopathic scoliosis (code range)
M41.30- M41.35	Thoracogenic scoliosis (code range)
M41.80- M41.87	Other forms of scoliosis (code range)
M41.9	Scoliosis, unspecified
M43.6	Torticollis
M48.00- M48.08	Spinal stenosis (code range)
M50.10- M50.13	Cervical disc disorder with radiculopathy (code range)

## Medical Policy: Evoked Potentials

Policy Number: 02.01.27

Page: 10 of 13

Code	Description
M50.20- M50.23	Other cervical displacement (code range)
M53.0	Cervicocranial syndrome
M53.1	Cervicobrachial syndrome
M53.80- M53.88	Other specified dorsopathies (code range)
M54.11	Radiculopathy, occipito-atlanto-axial region
M54.13	Radiculopathy, cervicothoracic region
M54.81	Occipital neuralgia
M96.3	Postlaminectomy kyphosis
M96.4	Postsurgical lordosis
M99.20- M99.29	Subluxation stenosis of neural canal (code range)
M99.30- M99.39	Osseous stenosis of neural canal (code range)
M99.40- M99.49	Connective tissue stenosis of neural canal (code range)
M99.50- M99.59	Intervertebral disc stenosis of neural canal (code range)
M99.60- M99.69	Osseous and subluxation stenosis of intervertebral foramina (code range)
M99.70- M99.79	Connective tissue and disc stenosis of intervertebral foramina (code range)

## REFERENCES

American Academy of Neurology. [Internet]. Cervical and ocular vestibular evoked myogenic potential testing: practice guidelines. 2017 [updated 2024 Feb 10; accessed 2026 Mar 30]. Available from: <https://www.aan.com/Guidelines/home/GuidelineDetail/877>

American Academy of Otolaryngology—Head and Neck Surgery. [Internet] Position statement: vestibular myogenic potential (VEMP) testing. [Updated 2021 Apr 21; accessed 2026 Mar 30]. Available from: <https://www.entnet.org/resource/position-statement-vestibular-myogenic-potential-vemp-testing/>

Barczyński M, et al. Randomized clinical trial of visualization versus neuromonitoring of recurrent laryngeal nerves during thyroidectomy. Br J Surg. 2009 Mar;96(3):240-6.

## **Medical Policy: Evoked Potentials**

**Policy Number: 02.01.27**

**Page: 11 of 13**

Cirocchi R, et al. Intraoperative neuromonitoring versus visual nerve identification for prevention of recurrent laryngeal nerve injury in adults undergoing thyroid surgery. *Cochrane Database Syst Rev*. 2019 Jan 19;1(1):CD012483.

Cohen H, et al. Relationship between clinical measures of hearing and clinical measures of vestibular function. *Am J Otolaryngol*. 2024 Jan-Feb;45(1):104052.

Cozzi AT, et al. Intraoperative neuromonitoring does not reduce the risk of temporary and definitive recurrent laryngeal nerve damage during thyroid surgery: a systematic review and meta-analysis of endoscopic findings from 73,325 nerves at risk. *J Pers Med*. 2023 Sep 23;13(10):1429.

Cui W, et al. Assessment of vestibular-evoked myogenic potentials in Parkinson's disease: a systematic review and meta-analysis. *Brain Sci*. 2022 Jul 21;12(7):956.

Di Martino A, et al. Should evoked potential monitoring be used in degenerative cervical spine surgery? A systematic review. *J Orthop Traumatol*. 2019 Apr 2;20(1):19.

Diercks GR, et al. Pediatric intraoperative nerve monitoring during thyroid surgery: A review from the American Head and Neck Society Endocrine Surgery Section and the International Neural Monitoring Study Group. *Head Neck*. 2022 Jun;44(6):1468-1480.

Evans AB. [Internet] Clinical utility of evoked potentials. [Updated 2019 Oct 25; accessed 2026 Mar 30]. Available from: <https://emedicine.medscape.com/article/1137451-overview>

Fife TD, et al. Practice guideline: Cervical and ocular vestibular evoked myogenic potential testing: report of the guideline development, dissemination, and implementation subcommittee of the American Academy of Neurology. *Neurology*. 2017 Nov 28;89(22):2288-2296.

Gronseth GS, et al. Practice parameter: the usefulness of evoked potentials in identifying clinically silent lesions in patients with suspected multiple sclerosis (an evidence-based review): report of the quality standards subcommittee of the American Academy of Neurology. *Neurology*. 2000 May 9;54(9):1720-5.

Jha MK, et al. Visual Evoked Potentials in Primary Open Angle Glaucoma. *J Neurodegener Dis*. 2017;2017:9540609. Epub 2017 Jul 20.

Mousa MF, et al. Evaluation of hemifield sector analysis protocol in multifocal visual evoked potential objective perimetry for the diagnosis and early detection of glaucomatous field defects. *Korean J Ophthalmol*. 2014 Feb;28(1):49-65. Epub 2014 Jan 21.

Muengtaweepongsa S. [Internet] General principles of somatosensory evoked potentials. Overview. [updated 2024 Oct 16; accessed 2026 Mar 30]. Available from: [https://emedicine.medscape.com/article/1139906-overview?&icd=login\\_success\\_email\\_match\\_fpf](https://emedicine.medscape.com/article/1139906-overview?&icd=login_success_email_match_fpf)

Pillai C, et al. Sensitivity and specificity of short-duration transient visual evoked potentials (SD-tVEP) in discriminating normal from glaucomatous eyes. *Invest Ophthalmol Vis Sci*. 2013 Apr 23;54(4):2847-52.

## Medical Policy: Evoked Potentials

Policy Number: 02.01.27

Page: 12 of 13

Verrecchia L, et al. Diagnostic accuracy of ocular vestibular evoked myogenic potentials for superior canal dehiscence syndrome in a large cohort of dizzy patients. *Ear Hear.* 2019 Mar/Apr;40(2):287-294.

Waisbourd M, et al. Short-duration transient visual evoked potentials and color reflectivity discretization analysis in glaucoma patients and suspects. *Int J Ophthalmol.* 2017 Feb 18;10(2):254-261.

### SEARCH TERMS

SEEPs

### CENTERS FOR MEDICARE AND MEDICAID SERVICES (CMS)

[NCD - Evoked Response Tests \(160.10\)](#) [accessed 2026 Mar 30]

[LCD - Visual Electrophysiology Testing \(L36831\)](#) [accessed 2026 Mar 30]

### PRODUCT DISCLAIMER

- Services are contract dependent; if a product does not cover a service, medical policy criteria do not apply.
- If a commercial product (including an Essential Plan or Child Health Plus product) covers a specific service, medical policy criteria apply to the benefit.
- If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit.
- If a Medicare product (including Medicare HMO-Dual Special Needs Program (DSNP) product) covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.
- If a Medicare HMO-Dual Special Needs Program (DSNP) product DOES NOT cover a specific service, please refer to the Medicaid Product coverage line.

### POLICY HISTORY/REVISION

#### Committee Approval Dates

10/19/00, 12/20/01, 12/19/02, 04/20/06, 03/15/07, 03/20/08, 02/19/09, 02/18/10, 02/17/11, 02/16/12, 02/21/13, 02/20/14, 01/22/15, 01/21/16, 01/19/17, 01/18/18, 03/21/19, 03/19/20, 03/18/21, 05/19/22, 05/18/23, 05/16/24, 05/22/25, 05/21/26

Date	Summary of Changes
05/21/26	<ul style="list-style-type: none"><li>• Annual review; policy intent unchanged</li></ul>
05/22/25	<ul style="list-style-type: none"><li>• Annual review, policy intent unchanged.</li></ul>

**Medical Policy: Evoked Potentials**

**Policy Number: 02.01.27**

**Page: 13 of 13**

01/01/25	<ul style="list-style-type: none"><li>• Summary of changes tracking implemented.</li></ul>
10/19/00	<ul style="list-style-type: none"><li>• Original effective date</li></ul>