

MEDICAL POLICY

| MEDICAL POLICY DETAILS | |
|-------------------------|---|
| Medical Policy Title | Inflammatory Markers of Coronary Artery Disease Risk |
| Policy Number | 2.02.15 |
| Category | Technology Assessment |
| Original Effective Date | 12/18/02 |
| Committee Approval Date | 05/21/03, 05/19/04, 03/17/05, 01/19/06, 11/16/06, 09/20/07, 09/18/08, 02/19/09, 03/18/10, 04/21/11, 04/19/12, 04/18/13, 04/17/14, 04/16/15, 06/16/16, 07/20/17, 06/21/18, 06/20/19, 06/18/20, 06/17/21, 06/16/22, 07/20/23, 07/18/24 |
| Current Effective Date | 07/18/24 |
| Archived Date | N/A |
| Archive Review Date | N/A |
| Product Disclaimer | <ul style="list-style-type: none"> • <i>Services are contract dependent; if a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</i> • <i>If a commercial product (including an Essential Plan or Child Health Plus product), medical policy criteria apply to the benefit.</i> • <i>If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit.</i> • <i>If a Medicare product (including Medicare HMO-Dual Special Needs Program (DSNP) product) covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</i> • <i>If a Medicare HMO-Dual Special Needs Program (DSNP) product DOES NOT cover a specific service, please refer to the Medicaid Product coverage line.</i> |

POLICY STATEMENT

- I. Based upon our criteria and assessment of the peer-reviewed literature, including the January 2003 recommendation put forth by the American Heart Association and Centers for Disease Control and Prevention, the use of high sensitivity C-reactive protein (hs-CRP) testing for primary prevention in the clinical setting is considered **medically appropriate** for those individuals who are at intermediate risk (10%-20%) of heart disease over the next 10 years by conventional risk scoring (e.g., Framingham Heart Study criteria) and who are free of non-cardiac conditions that are known to increase CRP (e.g., rheumatoid arthritis, chronic inflammatory processes).
- II. Based upon our criteria and assessment of the peer-reviewed literature, all other indications for hs-CRP testing, aside from the indication above, are considered **not medically necessary**.
- III. Based upon our criteria and assessment of the peer-reviewed literature, measurement of other inflammatory markers including but not limited to lipoprotein-associated phospholipase A₂ (Lp-PLA₂) and plasma myeloperoxidase (MPO) in the assessment of cardiovascular risk by any method has not been proven to improve health outcomes and, therefore, is considered **investigational**.

Refer to Corporate Medical Policy #11.01.03 Experimental or Investigational Services

POLICY GUIDELINES

- I. To be eligible for coverage of hs-CRP testing, a patient must be categorized as at a 10 to 20% higher risk (intermediate risk) than the average individual. Determination of increased risk is based on the Framingham Heart Study that identified patients who can be classified as either low, intermediate, or high risk for the cardiovascular events in the next 10 years. The classification is based on factors such as high blood pressure, high blood cholesterol, smoking, obesity, diabetes, and physical inactivity.

Medical Policy: INFLAMMATORY MARKERS OF CORONARY ARTERY DISEASE RISK

Policy Number: 2.02.15

Page: 2 of 9

- II. Several of the hs-CRP tests have received 510(k) marketing clearance from the U.S. Food and Drug Administration (FDA). In 2003, the FDA cleared for marketing an enzyme linked immunoabsorbent (ELISA) test, the PLAC test (diaDexus, San Francisco, CA) to measure levels of Lp-PLA₂.

DESCRIPTION

High sensitivity C-reactive protein (hs-CRP) is a nonspecific, acute-phase reactant, produced by the liver as a marker of inflammatory processes. Traditionally CRP has been used to monitor inflammatory processes, such as infections or autoimmune diseases. Chronic inflammatory disorders, including autoimmune diseases and malignancies can produce persistent increases in serum CRP concentrations. Studies suggest the association of low-level chronic inflammation during atherogenesis. The use of technologies collectively known as hs-CRP, including enzyme linked immunoabsorbent assays (ELISA) and other techniques using monoclonal antibodies, has allowed for a greater precision in detecting the lower levels of CRP that are related to chronic inflammation in otherwise healthy individuals. Results from studies indicate a correlation between hs-CRP levels and coronary artery disease. It is theorized that the increased sensitivity of an hs-CRP test should be able to detect that activity as a marker for cardiovascular disease, either current or future.

Lipoprotein-associated phospholipase A₂ (Lp-PLA₂), also known as platelet-activating factor acetylhydrolase, is an enzyme that hydrolyses phospholipids and is primarily associated with low-density lipoproteins. Accumulating evidence has suggested that Lp-PLA₂ is a biomarker of coronary artery disease and may have a proinflammatory role in the progression of atherosclerosis. The recognition that atherosclerosis represents, in part, an inflammatory process has created considerable interest in measurement of proinflammatory factors as part of cardiovascular disease risk assessment. Large, prospective studies are needed to establish whether measurement of Lp-PLA₂ biomarkers will be more predictive of cardiovascular disease (CVD) than conventional lipid risk factors.

Plasma myeloperoxidase (MPO), an abundant leukocyte enzyme, is elevated in culprit lesions that have fissured or ruptured in patients with sudden death from cardiac causes. Research suggests a mechanistic link between myeloperoxidase and both inflammation and cardiovascular disease risk. It has been proposed that elevated plasma MPO levels may be an independent predictor of endothelial dysfunction, angiographically-evident CAD and cardiac risk. There is a lack of scientific evidence regarding how measurements of MPO would affect management of individuals at risk for or patients with CVD. Large randomized controlled studies are needed to ascertain the clinical value of MPO in the management of CVD.

RATIONALE

Use of hs-CRP in Primary Prevention of Cardiovascular Disease

Several prospective epidemiologic studies have suggested that the measurement of hs-CRP may be an independent risk factor for cardiovascular disease.

Use of hs-CRP in Secondary Prevention of Cardiovascular Disease

Scientific evidence supports the theory that hs-CRP is a strong and independent marker for future heart events in patients who have already been assessed to be at a 10 to 20% greater risk than the average individual. Based on this information, use of the hs-CRP test to further evaluate this group of patients may result in a change in treatment and/or lifestyles that could decrease the risk for future cardiac events.

No clinical trials have been completed in which a population has been randomly allocated to hs-CRP screening compared with a control population group not allocated to hs-CRP screening with both groups followed up prospectively to determine the benefits and harms of the screening.

The American Heart Association (AHA) and Centers for Disease Control and Prevention (CDC) has issued the following recommendation regarding the role of hs-CRP measurements in clinical practice (2020): it is reasonable to measure hs-CRP as an adjunct to the major risk factors to further assess absolute risk for coronary disease primary prevention. At the discretion of the physician, the measurement is considered optional, based on the moderate level of evidence (Evidence Level C). In this role, hs-CRP measurement appears to be best employed to detect enhanced absolute risk in persons in whom multiple risk factor scoring projects a 10-year CHD risk in the range of 10% to 20% (Evidence Level B). However,

Medical Policy: INFLAMMATORY MARKERS OF CORONARY ARTERY DISEASE RISK

Policy Number: 2.02.15

Page: 3 of 9

the benefits of this strategy or any treatment based on this strategy remain uncertain. Individuals at low risk (10% per 10 years) will be unlikely to have a high risk (20%) identified through hs-CRP testing. Individuals at high risk (20% risk over 10 years) or with established atherosclerotic disease generally should be treated intensively regardless of their hs-CRP levels, so the utility of hs-CRP in secondary prevention appears to be more limited. In patients with stable coronary disease or acute coronary syndromes, hs-CRP measurement may be useful as an independent marker for assessing likelihood of recurrent events, including death, myocardial infarction, or restenosis after percutaneous coronary intervention. However, secondary preventive interventions with proven efficacy should not be dependent on hs-CRP levels. Further, serial testing of hs-CRP should not be used to monitor the effects of treatment.

Duprez et al. (2016) studied the association of GlycA and inflammatory biomarkers with future death and disease. A total of 6,523 subjects in the Multi-Ethnic Study of Atherosclerosis who were free of overt cardiovascular disease (CVD) and in generally good health had baseline blood analyzed for high-sensitivity C-reactive protein (hsCRP), interleukin-6 (IL-6), and d-dimer. A spectral deconvolution algorithm was used to quantify GlycA signal amplitudes from automated NMR LipoProfile test spectra. Median follow-up was 12.1 years. The authors found that relative risk per SD of GlycA, IL-6, and d-dimer for total death (n = 915); for total CVD (n = 922); and for chronic inflammatory-related severe hospitalization and death (ChIRd) (n = 1324) ranged from 1.05 to 1.20, independently of covariates. In contrast, prediction from hsCRP was statistically explained by adjustment for other inflammatory variables. Only GlycA was predictive for total cancer (n = 663). Women had 7% higher values of all inflammatory biomarkers than men and had a significantly lower GlycA prediction coefficient than men in predicting total cancer. The authors concluded that GlycA derived from NMR is associated with risk for total death, CVD, and chronic inflammatory-related severe hospitalization and death, and cancer. This novel biomarker reflecting the risk of death, CVD, ChIRd, and cancer may have the potential to improve risk assessment.

Lipoprotein-associated Phospholipase A₂ (Lp-PLA₂) as an Independent Biomarker

Current studies generally report the utility of Lp-PLA₂ as an independent biomarker for coronary artery disease and recurrent cardiac events. However, Lp-PLA₂ was not found to be an independent marker for subclinical atherosclerosis, and a study of the Atherosclerosis Risk in Communities (ARIC) cohort found that routine measurement of Lp-PLA₂ did not improve existing risk stratification models that use traditional risk factors. Interventional studies involving Lp-PLA₂ suggest that the level of Lp-PLA₂ is modifiable by antihyperlipidemics. An ad hoc study of the Pravastatin or Atorvastatin Evaluation and Infection Therapy: Thrombolysis in Myocardial Infarction 22 (PROVE IT-TIMI 22) trial concluded that the 30-day Lp-PLA₂ level was independently associated with an increased risk of cardiovascular events. Another ad hoc study from the Diabetes and Combined Lipid Therapy Regimen (DIACOR) trial demonstrated improved Lp-PLA₂ levels compared to baseline, with no difference found between treatment groups among 300 patients with diabetes and mixed dyslipidemias randomized to either fenofibrate, simvastatin, or both for 12 weeks.

Results of two large-scale observational studies have suggested that Lp-PLA₂ is an independent risk factor for coronary heart disease in men. However, the key outcome of cardiac risk assessment is an improvement in health outcomes. Improved risk prediction does not by itself result in improved health outcomes. To improve outcomes, clinicians must have the tools to translate this information into clinical practice. This requires guidelines that incorporate emerging risk factors into existing risk prediction models and have been demonstrated to classify patients into risk categories with greater accuracy. Predictive models also need to be accompanied by treatment guidelines that target intervention toward patients who will get the most benefit. At present, measurements of Lp-PLA₂ are not a component of the guidelines developed by the National Cholesterol Education Program Adult Treatment Panel III.

Zhang et al. (2021) performed a prospective study to investigate the association between cardiovascular disease and Lp-PLA₂. A total of 823 individuals at a high risk of stroke were screened and followed at three (3), six (6), 12, and 24 months. Among the 823 participants, 286 had varying degrees of carotid artery stenosis and 18 participants had cerebrovascular events. The level of Lp-PLA₂ was higher in the group with cerebrovascular events than in the group without cerebrovascular events (662.81 ± 111.25 vs 559.86 ± 130.05, p < 0.001). No statistical difference was found between the other parameters of the event group, such as HDL, LDL, and the no event group. The incidence of cerebrovascular events in the stenosis group was higher than that in the no stenosis group but no statistically significant difference was noted. The authors concluded that the level of Lp-PLA₂ was positively correlated with the degree of carotid artery stenosis and predicted cerebrovascular events. The study limitations included sample size; the study follow-

Medical Policy: INFLAMMATORY MARKERS OF CORONARY ARTERY DISEASE RISK

Policy Number: 2.02.15

Page: 4 of 9

up time was only two (2) years, and the number of cerebrovascular events that eventually occurred was relatively small. Also of note, the study was conducted at a single center and the study population mainly included people aged > 40 years at a high risk of stroke, so the results of the study only represented a small part of the population. Furthermore, the study did not address how integrating measurements of Lp-PLA₂ to clinical care alters patient management and improves clinical outcomes.

While studies have suggested that statin drugs and fibrates may reduce levels of Lp-PLA₂, it is not known whether such drug therapy in patients not already considered candidates, based on other well established risk factors would ultimately decrease the incidence of coronary heart disease. Although results of studies of Lp-PLA₂ test are promising, its biological role is not yet understood, its ability to improve on existing risk stratification methods is uncertain, and its clinical utility remains in question, particularly when compared to currently available methods for cardiovascular risk reduction. The extent to which antihyperlipidemics modify the level of Lp-PLA₂ beyond their established therapeutic use, and thereby altering cardiac outcomes, is unknown.

Risk Prediction for Stroke

While some studies have shown that levels of both Lp-PLA₂ and C-reactive protein were higher in stroke cases, improved risk prediction does not necessarily result in improved outcomes. Results of studies have not been incorporated into clinical management.

Several studies have assessed the value of MPO as a predictor of the risk of cardiovascular events in patients presenting with chest pain or acute coronary syndrome and chronic heart failure. MPO levels have also been evaluated as an inflammatory marker of future coronary artery disease (CAD) in apparently healthy individuals (Meuwesen, et al. 2007). Although studies of MPO testing indicate a possible relationship between elevated levels and cardiac risk, its ability to improve on existing risk stratification methods is unclear. (Roman, et al., 2008; Stefanescu, et al., 2008). Furthermore, in the studies evaluating MPO various methods of testing were used, making comparisons difficult. The body of evidence evaluating MPO as a potential cardiac biomarker is insufficient to support an increased predictive value as compared to traditional testing or for recommending medical management based on MPO values that would improve clinical outcomes.

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*
- *Code Key: Experimental/Investigational = (E/I), Not medically necessary/appropriate = (NMN).*

CPT Codes

| Code | Description |
|-------------|--|
| 83698 (E/I) | Lipoprotein-associated phospholipase A2, (Lp-PLA ₂) |
| 83876 (E/I) | Myeloperoxidase (MPO) |
| 86141 | C-reactive protein, high sensitivity, (hsCRP) |
| 0052U (E/I) | Lipoprotein, blood, high resolution fractionation and quantitation of lipoproteins, including all five major lipoprotein classes and subclasses of HDL, LDL, and VLDL by vertical auto profile ultracentrifugation (VAP Cholesterol Test, VAP Diagnostics Laboratory, Inc) |
| 0377U (E/I) | Cardiovascular disease, quantification of advanced serum or plasma lipoprotein profile, by nuclear magnetic resonance (NMR) spectrometry with report of a lipoprotein profile (including 23 variables) (Liposcale, CIMA Sciences, LLC) |

Medical Policy: INFLAMMATORY MARKERS OF CORONARY ARTERY DISEASE RISK**Policy Number: 2.02.15****Page: 5 of 9**

| Code | Description |
|-------------|--|
| 0415U (E/I) | Cardiovascular disease (acute coronary syndrome [ACS]), IL-16, FAS, FASLigand, HGF, CTACK, EOTAXIN, and MCP-3 by immunoassay combined with age, sex, family history, and personal history of diabetes, blood, algorithm reported as a 5-year (deleted risk) score for ACS (SmartHealth Vascular Dx, Morningstar Laboratories, LLC, SmartHealth DX) (<i>Effective 10/01/23</i>) |
| 0466U (E/I) | Cardiology (coronary artery disease [CAD]), DNA, genome-wide association studies (564856 single-nucleotide polymorphisms [SNPs], targeted variant genotyping), patient lifestyle and clinical data, buccal swab, algorithm reported as polygenic risk to acquired heart disease (<i>Effective 07/01/24</i>) |

Copyright © 2024 American Medical Association, Chicago, IL

HCPCS Codes

| Code | Description |
|---------------------|--------------------|
| No specific code(s) | |

ICD10 Codes

| Code | Description |
|---------------------|--|
| E71.30 | Disorder of fatty-acid metabolism, unspecified |
| E75.21 | Fabry (-Anderson) disease |
| E75.22 | Gaucher disease |
| E75.240- E75.249 | Niemann-Pick disease (code range) |
| E75.3 | Sphingolipidosis, unspecified |
| E75.5 | Other lipid storage disorders |
| E75.6 | Lipid storage disorder, unspecified |
| E77.0-E77.9 | Disorders of glycoprotein metabolism (code range) |
| E78.0-E78.9 | Disorders of lipoprotein metabolism and other lipidemias (code range) |
| E88.1 | Lipodystrophy, not elsewhere classified |
| E88.2 | Lipomatosis, not elsewhere classified |
| E88.89 | Other specified metabolic disorders |
| I20.0-I20.9 | Angina pectoris (code range) |
| I21.01-I22.1 | ST elevation (STEMI) myocardial infarction of anterior or inferior wall (code range) |
| I24.1 | Dressler's syndrome |
| I25.110-I25.119 | Atherosclerotic heart disease of native coronary artery with angina pectoris (code range) |
| I25.2 | Old myocardial infarction |
| I25.700-I25.799 | Atherosclerosis of coronary artery bypass graft(s) and coronary artery of transplanted heart with angina pectoris (code range) |

Medical Policy: INFLAMMATORY MARKERS OF CORONARY ARTERY DISEASE RISK**Policy Number: 2.02.15****Page: 6 of 9**

| Code | Description |
|-------------|--|
| I70.0 | Atherosclerosis of aorta |
| I70.1 | Atherosclerosis of renal artery |
| Z82.41 | Family history of sudden cardiac death |
| Z82.49 | Family history of ischemic heart disease and other diseases of the circulatory system |
| Z86.711 | Personal history of pulmonary embolism |
| Z86.718 | Personal history of other venous thrombosis and embolism |
| Z86.72 | Personal history of thrombophlebitis |
| Z86.73 | Personal history of transient ischemic attack (TIA), and cerebral infarction without residual deficits |
| Z86.74 | Personal history of sudden cardiac arrest |
| Z86.79 | Personal history of other diseases of the circulatory system |

REFERENCES

- *Abd TT, et al. The role of C-reactive protein as a risk predictor of coronary atherosclerosis: implications from the JUPITER trial. Curr Atheroscler Rep 2011 Apr;13(2):154-61.
- Adam LN, et al. Genetic and clinical study of myeloperoxidase's association with coronary artery disease. The Egyptian Heart Journal 2024;76:27.
- *Alvarez GB, et al. High-sensitivity C-reactive protein in high-grade carotid stenosis: risk marker for unstable carotid plaque. J Vasc Surg 2003 Nov;38(5):1018-24.
- *Aviles RJ, et al. Inflammation as a risk for atrial fibrillation Circ 2003 Dec 16;108(24):3006-10.
- *Baldus S, et al. Myeloperoxidase serum levels predict risk in patients with acute coronary syndromes. Circ 2003 Sep 23;108(12):1440-5.
- *Ballantyne CM, et al. Lipoprotein-associated phospholipase A2, high-sensitivity C-reactive protein, and risk for incident coronary heart disease in middle-aged men and women in the atherosclerosis risk in communities (ARIC) study. Circ 2004 Feb 24;109(7):837-42.
- *Bhatti S, et al. Lp-PLA2 as a marker of cardiovascular diseases. Curr Atheroscler Rep 2010;12:140-4.
- *Blake GJ, et al. A prospective evaluation of lipoprotein-associated phospholipase A(2) levels and the risk of future cardiovascular events in women. J Am Coll Cardiol 2001 Nov;38(5):1302-6.
- *Brennan M, et al. Prognostic value of myeloperoxidase in patients with chest pain. N Engl J Med 2003 Oct 23;349(17):1595-604.
- *Chang C, et al. The myeloperoxidase -463G/A polymorphism and coronary artery disease risk: A meta-analysis of 1938 cases and 1990 controls. Clin Biochem 2013;46(16-17):1644-8.
- Chong Y, et al. Biomarkers for prediction of cardiovascular events in community-dwelling adults aged 40 or older. Int Heart J 2020 Jan 31;61(1):109-114.
- *Cook NR, et al. Clinical utility of lipoprotein-associated phospholipase A2 for cardiovascular disease prediction in a multiethnic cohort of women. Clin Chem 2012 Sep;58(9):1352-63.
- DeStefano A. et al. Lp-PLA2, a new biomarker of vascular disorders in metabolic diseases. Int J Immunopathol Pharmacol 2019 Jan-Dec;33:2058738419827.

Medical Policy: INFLAMMATORY MARKERS OF CORONARY ARTERY DISEASE RISK

Policy Number: 2.02.15

Page: 7 of 9

- *Duprez DA, et al. High-density lipoprotein subclasses and noncardiovascular, noncancer chronic inflammatory-related events versus cardiovascular events: the multi-ethnic study of atherosclerosis. J Am Heart Assoc 2015 Sep 14;4(9):e002295.
- *Elkind MS, et al. High-sensitivity C-reactive protein, lipoprotein-associated phospholipase A2, and outcome after ischemic stroke. Arch Intern Med 2006 Oct 23;166(19):2073-80.
- *Epps KC, et al. Lp-PLA2 – a novel risk factor for high-risk coronary and carotid artery disease. J Intern Med 2011;269:94–106.
- Fatemi S, et al. Lp-PLA2 activity and mass and CRP are associated with incident symptomatic peripheral arterial disease. Sci Rep 2019;9:5609.
- *Folsom AR, et al. An assessment of incremental coronary risk prediction using C-reactive protein and other novel risk markers; the atherosclerosis risk in communities study. Arch Intern Med 2006 Jul 10;166(13):1368-73.
- *Grundy SM, Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. Circ 2004 Jul 13;110(2):227-39.
- *Grundy SM, et al. Prevention conference V beyond secondary prevention: identifying the high-risk patient for primary prevention: medical office assessment: writing group I. Circ 2000;1010:e3.
- *Hatoum IJ, et al. Lipoprotein-associated phospholipase A2 activity improves risk discrimination of incident coronary heart disease among women. Am Heart J 2011;161:516-22.
- Hu Y, et al. Lipoprotein-associated phospholipase A2 is a risk factor for diabetic kidney disease. Diabetes Res Clin Pract 2019 Mar 20;150:194-201.
- *Kardys I, et al. Lipoprotein-associated phospholipase A2 and measures of extra atherosclerosis: the Rotterdam Study. Arterioscler Thromb Vasc Biol 2006 Mar;26(3):631-6.
- *Koenig W, et al. C-reactive protein modulates risk prediction based on the Framingham Score: implications for future risk assessment: results from a large cohort study in southern Germany. Circ 2004 Mar 23;109(11):1349-53.
- *Koenig W, et al. Lipoprotein-associated phospholipase A2 adds to risk prediction of incident coronary events by C-reactive protein in apparently healthy middle-aged men from the general population: results from the 14-year follow-up of a large cohort from southern Germany. Circ 2004 Oct 5;110(14):1903-8.
- *Koenig W, et al. Lipoprotein-associated phospholipase A2 predicts future cardiovascular events in patients with coronary heart disease independently of traditional risk factors, markers of inflammation, renal function, and hemodynamic stress. Arterioscler Thromb Vasc Biol 2006 Jul;26(7):1586-93.
- *Koenig W. High-sensitivity C-reactive protein and atherosclerotic disease: from improved risk prediction to risk-guided therapy. Int J Cardiol 2013 Oct 15;168(6):5126-34.
- *Lin J, et al. Association of Lp-PLA2-A and early recurrence of vascular events after TIA and minor stroke. Neurology 2015 Nov 3;85(18):1585-91.
- Ling Y, et al. Relationship between plasma lipoprotein-associated phospholipase A2 concentrations and apolipoprotein in stable coronary artery disease patients. Dis Markers 2020 Sep 24;2020:8818358.
- *Liu J, et al. Association between the lipoprotein-associated phospholipase A2 activity and the progression of subclinical atherosclerosis. J Atheroscler Thromb 2014;21(6):532-42.
- Maestrini I, et al. Analysis of the association of MPO and MMP-9 with stroke severity and outcome. Neurology 2020 Jul 7;95(1):e97-e108.
- *Maiolino G, et al. Lipoprotein-associated phospholipase a2 prognostic role in atherosclerotic complications. World J Cardiol 2015 Oct 26;7(10):609-20.

Medical Policy: INFLAMMATORY MARKERS OF CORONARY ARTERY DISEASE RISK

Policy Number: 2.02.15

Page: 8 of 9

Mahat RK, et al. Association of myeloperoxidase with cardiovascular disease risk factors in prediabetic subjects. Diabetes Metab Syndr 2019 Jan – Feb;13(1):396-400.

*Mangili A, et al. Lipoprotein-associated phospholipase a2, a novel cardiovascular inflammatory marker, in HIV-infected patients. Clin Infect Dis 2014 Mar;58(6):893-900.

*Muhlestein JB, et al. The reduction of inflammatory biomarkers by statin, fibrate, and d combination therapy among diabetic patients with mixed dyslipidemia: the DIACOR (Diabetes and Combined Lipid Therapy Regimen) study. J Am Coll Cardiol 2006 Jul 18;48(2):396-401.

Nadel J, et al. Arterial myeloperoxidase in the detection and treatment of vulnerable atherosclerotic plaque: a new dawn for an old light. European Society of Cardiology Cardiovascular Research 2023;119:112-120.

*Ndrepepa G. Myeloperoxidase – a bridge linking inflammation and oxidative stress with cardiovascular disease. Clinica Chimica Acta 2019 Jun;493:36-51.

*O'Donoghue M, et al. Lipoprotein-associated phospholipase A2 and its association with cardiovascular outcomes in patients with acute coronary syndromes in the PROVE IT-TIMI 22 (PRavastin Or atorVastin Evaluation and Infection Therapy-Thrombolysis In Myocardial Infarction) trial. Circ 2006 Apr 11;113(14):1745-52.

*Pai JK, et al. Inflammatory markers and the risk of coronary heart disease in men and women. NEJM 2004 Dec 16;351(25):2599-610.

*Pearson TA, et al. Markers of inflammation and cardiovascular disease: application to clinical and public health practice: a statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. Circ 2003 Jan 23;107(3):499-511.

Ramachandra CJA, et al. Myeloperoxidase as a multifaceted target for cardiovascular protection. Antioxid Redox Signal 2020 May 20;32(15):1135-1149.

*Roman RM, et al. Myeloperoxidase and coronary arterial disease: from research to clinical practice. Arq Bras Cardiol 2008 Jul;91(1):e11-9.

*Stefanescu A, et al. Prognostic value of plasma myeloperoxidase concentration in patients with stable coronary artery disease. Am Heart J 2008 Feb;155(2):356-60.

Sun L, et al. Causal effect of Lipoprotein-associated phospholipase A2 activity on coronary artery disease and myocardial infarction: a two-sample Mendelian randomization study. Clinica Chimica Acta 2021;523:491-496.

Tan Y, et al. Addition of plasma myeloperoxidase and trimethylamine n-oxide to the GRACE score improves prediction of near-term major adverse cardiovascular events in patients with ST-segment elevation myocardial infarction. Frontiers in Pharmacology Sept 2021;12:Article#632075.

*Tang WH, et al. Plasma myeloperoxidase levels in patients with chronic heart failure. Am J Cardiol 2006 Sep 15;98(6):796-9.

*Vasan RS, et al. Inflammatory markers and risk of heart failure in elderly subjects without prior myocardial infarction: the Framingham Heart Study. Circ 2003 Mar 25;107(11):1486-91.

*Yeh ETH, et al. Coming of age of C-reactive protein: using inflammation markers in cardiology. Circ 2003 Jan 28;107(3):370-2.

Zhang F, et al. Lp-PLA2 evaluates the severity of carotid artery stenosis and predicts the occurrence of cerebrovascular events in high stroke-risk populations. J Clin Lab Anal 2021 Mar;35(3): e23691.

Zhang H, et al. The relationship of lipoprotein-associated phospholipase A2 activity with the seriousness of coronary artery disease. BMC Cardiovasc Disord 2020;20:295.

*Key Article

Medical Policy: INFLAMMATORY MARKERS OF CORONARY ARTERY DISEASE RISK

Policy Number: 2.02.15

Page: 9 of 9

KEY WORDS

Cardiac disease risk, CRP, hs-CRP, Lp-PLA2, PLAC test, plasma myeloperoxidase (MPO).

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

Based upon our review, High Sensitivity C-Reactive Protein Testing (hsCRP) and Lipid Testing is not addressed in National or Regional Medicare coverage determinations or policies.

However, please refer to the Medicare Managed Care Manual/Chapter 4: Benefits and Beneficiary Protections (Rev.121, Issued: 04-22-16)/Section 90 National and Local Coverage Determinations/Subsection 90.4.1 MAC with Exclusive Jurisdiction over a Medicare Item or Service:

In some instances, one Medicare A/B MAC processes all of the claims for a particular Medicare-covered item or service for all Medicare beneficiaries around the country. This generally occurs when there is only one provider of a particular item or service (for example, certain pathology and lab tests furnished by independent laboratories). In this situation, MA plans must follow the coverage policy reflected in an LCD issued by the A/B MAC that enrolled the provider and processes all the Medicare claims for that item or service.

[<https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Internet-Only-Manuals-IOMs-Items/CMS019326>] accessed 06/06/24.