

MEDICAL POLICY

| | |
|------------------------|------------------------|
| Medical Policy Title | Kidney Transplantation |
| Policy Number | 07.02.04 |
| Current Effective Date | April 16, 2026 |
| Next Review Date | April 2027 |

Our medical policies are guides to evaluate technologies or services for medical necessity. Criteria are established through the assessment of evidence based, peer-reviewed scientific literature, and national professional guidelines. Federal and state law(s), regulatory mandates and the member's subscriber contract language are considered first in the determination of a covered service.

(Link to [Product Disclaimer](#))

POLICY STATEMENT(S)

- I. Kidney transplants are **medically appropriate** for candidates with documented progressive end stage renal disease (ESRD) when **EITHER** of the following indications are present:
 - A. A measured (actual urinary collection) creatinine clearance level or calculated glomerular filtration rate (GFR) (or another reliable formula) less than or equal to 20ml/min; **or**
 - B. The initiation of dialysis.
- II. Kidney retransplants after a failed primary kidney transplant are considered **medically necessary** when the criteria in Policy Statement I are met.
- III. The following conditions are absolute contraindications to kidney transplantation:
 - A. Presence of malignancy (other than non-melanoma skin cancers), unless malignancy has been completely resected, or unless (upon medical review) it is determined that malignancy has been treated with a small likelihood of recurrence and acceptable future risks;
 - B. Ongoing or recurring infections that are not effectively treated;
 - C. Serious cardiac or pulmonary, other ongoing comorbid conditions that create an inability to tolerate transplant surgery;
 - D. Demonstrated non-adherence to medical recommendations, which places the organ at risk.
- IV. Kidney transplantation is considered a relative contraindication in human immunodeficiency virus (HIV) positive individuals, unless **ALL** of the following criteria are met:
 - A. Cluster of Differentiation 4 (CD4) count greater than 200 cells/mm³;
 - B. Undetectable HIV-1 ribonucleic acid (RNA);
 - C. On Stable anti-retroviral therapy for greater than three (3) months; **and**
 - D. All other criteria within the policy are met.

RELATED POLICIES

[Corporate Medical Policy](#)

Medical Policy: Kidney Transplantation

Policy Number: 07.02.04

Page: 2 of 10

07.02.01 Pancreas Transplant: (Pancreas Transplant Alone, Pancreas Transplant after Kidney Transplant, Simultaneous Pancreas Kidney Transplant): Islet Cell Transplant

POLICY GUIDELINE(S)

- I. Recipient Selection
 - A. Each individual considered for renal transplantation will have an evaluation completed by the transplant center for potential difficulties that would complicate and diminish the success of transplantation. Consideration will be given to the individual's risk of death without transplantation, along with the presence and severity of potential contraindications to transplantation. Candidates considered for transplant must be psychologically stable, demonstrate motivation and compliance, and have no ongoing problems with drug or alcohol abuse.
 - B. Nutritional issues are important predictors of surgical outcomes. For candidates with a body mass index (BMI) outside of the "normal range," documentation of dietary counseling will be required at the time of evaluation and while on the waiting list.
 - C. Diabetic complications often fall into the realm of relative contraindications (except for significant cardiovascular disease); however, renal transplantation is associated with improved survival in individuals with end-stage renal disease (ESRD) caused by type 1 diabetes mellitus. Individuals with diabetes may be candidates for combined kidney-pancreas transplantation.
- II. Prior authorization requirements are contract dependent. Approval for all transplants, including arrangements with an approved transplant center, may be required.
- III. Pre-transplant evaluation documentation must include the following clinical information (if testing is unable to be performed, the rationale for not performing the testing must be included in the documentation):
 - A. Clinical Evaluation:
 1. Confirmation of diagnosis;
 2. Identification of comorbidities;
 3. Current assessment of co-morbidities;
 4. Management of co-morbidities; and
 5. Consult notes (if applicable).
 - B. Psycho-Social Evaluation:
 1. Identification of stressors (family support, noncompliance issues, motivational issues, alcohol, or smoking/substance abuse).
 - C. Performance Status:
 1. Karnofsky performance score;

Medical Policy: Kidney Transplantation

Policy Number: 07.02.04

Page: 3 of 10

2. Palliative Performance Scale (PPS) score;
 3. Eastern Cooperative Oncology Group (ECOG) performance status; or
 4. Lansky Play-Performance Scale (for age 1 to 16 years).
- D. Oral Health Evaluation;
- E. Lab Tests:
1. Complete Blood Count (CBC), metabolic profile;
 2. Serologies: Cytomegalovirus (CMV); Hepatitis B and C; and
 3. Human Immunodeficiency Virus (HIV) Testing.
- F. Cardiac Assessment:
1. 12 Lead Electrocardiogram (EKG);
 2. Stress (exercise, nuclear, or dobutamine); and
 3. Echocardiogram (Echo) or Multigated Acquisition (MUGA) Scan.
- G. Pulmonary Assessment:
1. Chest x-ray;
 2. Pulmonary function tests (PFTs) for high-risk respiratory failure (COPD, emphysema, alpha 1-antitrypsin deficiency, hepatopulmonary syndrome, or significant smoking history); and
- H. Age-appropriate Screening Tests: Please refer to the U.S Preventive Services Task Force (USPSTF) [Internet] for list of age-appropriate screening guidelines (e.g., colorectal cancer screening, cervical cancer screening for guidance). [accessed 2026 Feb 25]. Available from: <https://uspreventiveservicestaskforce.org/uspstf/>
- IV. Hepatitis C virus (HCV) infections are common among individuals with chronic renal failure and result in significant morbidity and mortality. Therefore, the assessment of hepatitis C virus infection in the potential recipient has a major clinical significance. HCV infections are associated with an increased risk of death, irrespective of whether the individual stays on dialysis or has a renal transplant. Transplantation has a beneficial, rather than adverse, effect on long-term survival in an anti-HCV-positive individual. An HCV-positive status is not a contradiction for renal transplantation.
- V. A person who gives consent to be a live organ donor should be competent, willing to donate, free from coercion, medically and psychologically suitable, fully informed of the risks and benefits as a donor, and fully informed of the risks, benefits, and alternative treatment available to the recipient. The benefits to both donor and recipient must outweigh the risks associated with the donation and transplantation for the living donor organ.
- VI. Candidates may be wait-listed at more than one transplant center. Since waiting time priority is first calculated among candidates at all hospitals within the local donation area, listing at

Medical Policy: Kidney Transplantation

Policy Number: 07.02.04

Page: 4 of 10

transplant centers in different local allocation areas is recommended. Requirements for multiple-listed candidates may vary among transplant centers. When possible, results of tests used in the evaluation for the transplant at one center should be used at subsequent centers where the individual is listed.

VII. Transplant re-authorization must be completed annually while actively waiting for a transplant. Re-authorization documentation must be within the past 11 months (unless specified) and include the following clinical information (if testing is unable to be performed, the rationale must be included in the documentation). If the individual's health condition has not changed from the previous year some testing would not be applicable.

A. Clinical Evaluation:

1. Updated list of diagnoses to include identification of comorbidities, current assessment, and treatment plan.
2. Specialty consultation notes (if applicable).

B. Current functional ability as evidenced by current Karnofsky performance score (KPS); Palliative Performance Scale (PPS) score; ECOG, or Lansky Play-Performance Scale.

C. Follow-up Oral Health Evaluation:

D. Lab Tests:

1. CBC, metabolic profile;
2. Serologies: CMV, Hepatitis B and C; and
3. HIV testing (If applicable).

E. Cardiac Assessment:

1. 12 Lead EKG (If applicable);
2. Stress (exercise, nuclear, or dobutamine) (If applicable); and
3. Echo or Muga scan (If applicable).

F. Pulmonary Assessment:

1. Chest x-ray (If applicable);
2. Pulmonary function tests (PFTs) for high-risk respiratory failure (COPD, emphysema, α -1-antitrypsin deficiency, hepatopulmonary syndrome, or significant smoking history);
and

G. Age-appropriate Screening Tests: Please refer to the USPSTF [Internet] for a list of age-appropriate screening guidelines (e.g., colorectal cancer screening, cervical cancer screening for guidance). [accessed 2026 Feb 25]. Available from:

<https://uspreventiveservicestaskforce.org/uspstf/>

| |
|--------------------|
| DESCRIPTION |
|--------------------|

Medical Policy: Kidney Transplantation

Policy Number: 07.02.04

Page: 5 of 10

The kidneys are two bean-shaped organs located on each side of the spine just below the rib cage. Kidneys are about the size of a fist. Their main function is to filter blood toxins and transform the waste into urine to be eliminated. When kidneys lose filtering abilities, harmful levels of fluid and waste accumulate in the body, which can result in kidney (renal) failure. Estimated glomerular filtration rate (eGFR) is a calculation used to estimate how well the kidneys are filtering certain products produced by the body. In adults, the normal eGFR number is usually 90-100. Which means that the kidney function is 90-100%. eGFR declines with age, even in people without kidney disease.

The National Kidney Foundation (NKF) states that Stage 5 chronic kidney disease (end-stage renal disease) is eGFR less than 15 for 3 months or more, or the individual is on dialysis. GFR estimate is considered the best measure of kidney function (NKF 2023).

Individuals with kidney failure are treated with dialysis, or with transplant.

A kidney transplant is the removal of a healthy kidney from a living donor or deceased (cadaveric) donor and the placement of the kidney into a person whose kidneys are no longer functioning properly. Transplant recipients require life-long immunosuppression to prevent rejection. Individuals are prioritized for transplant by mortality risk and severity of illness criteria developed by the Organ Procurement and Transplantation Network (OPTN) and United Network of Organ Sharing (UNOS). Data collected in 2023 (Lentine et al) show over 28,000 kidney transplantations performed in the United States in that year, with a rise noted due to an increase in deceased donor transplants. Five-year graft survival rates were 90% for living donor transplants and 82.2% for deceased donor transplants (ages 18-34 years). There were more than 140,000 adult candidates on the waiting list in 2023.

SUPPORTIVE LITERATURE

Chaudhry et al (2022) published a systematic review that compared survival for waitlisted patients with kidney failure who received a transplant compared to those who remained on the transplant waiting list. A total of 48 observational studies were included in the systematic review, of which 18 studies were suitable for meta-analysis. Results demonstrated a 55% reduction in the risk of mortality in patients who received a transplant compared to those who remained on dialysis.

Kainz et al (2022), retrospectively compared the association of time on the waitlist with survival of those who received a second transplant to those who remained on the waitlist. A total of 2,346 individuals from the Austrian Dialysis and Transplant Registry and Eurotransplant were included in the study. Inclusion criteria were individuals with a failed first graft, who were over the age of 18 years, and waitlisted for a second kidney transplantation in Austria between the years of 1980 and 2019. Study results demonstrated that second kidney transplantation improved survival at 10 years of follow-up compared with those remaining on the waitlist. The survival difference diminished in the individuals with longer wait times after loss of the first allograft. Individuals with a waitlist for retransplantation of 1 year or less, gained 8.0 life months compared to those who remained on the waiting list for 8 years (0.1 life month gained).

Kidney Retransplant

Pediatric kidney transplant recipients will likely require a retransplant in their lifetime. Gupta et al

Medical Policy: Kidney Transplantation

Policy Number: 07.02.04

Page: 6 of 10

(2015) retrospectively analyzed OPTN data, focusing on patients who had an initial kidney transplant as children. A total of 2,281 patients were identified. Included were those who had their first transplant before 18 years of age and a second kidney transplant at any age following. In multivariate analysis, the length of first graft survival and age at second graft were significantly associated with second graft survival. Specifically, the first graft survival time of more than 5 years was associated with better second graft survival. However, patients who were between 15 and 20 years old at second transplant were at increased risk of second kidney graft failure compared with individuals in other age groups. It is likely that most pediatric renal allograft recipients will require one or more retransplants during their lifetime. Unfortunately, the use of a second or subsequent graft in pediatric recipients has inferior long-term graft survival rates compared with initial grafts, with decreasing rates with each subsequent graft (Saeed et al 2024).

More recent research is aimed at strategies to prevent transplant rejection, while cell-based therapies are being investigated to minimize the impacts of immunosuppression.

PROFESSIONAL GUIDELINE(S)

The 2021 Chronic Kidney Disease Epidemiology Collaboration (2021 CKD-EPI) removed race as a variable from the GFR equation and recalibrated the other coefficients variables (e.g., age, sex, and serum creatinine). Subsequently, the National Kidney Foundation and American Society of Nephrology Task Force recommended that the 2021 CKD-EPI equation be implemented for eGFR reporting (Delgado et al 2022).

OPTN requires all transplant hospitals to use race-neutral calculations when estimating a candidate's GFR for any purpose covered by OPTN policy. Kidney transplant programs are required to assess their waiting lists and correct wait times for any African American kidney candidates disadvantaged by having their kidney function overestimated because of the use of a race-inclusive calculation (OPTN 2023).

Kidney Disease Improving Global Outcomes (KDIGO) 2024 Clinical Practice Guideline for the Evaluation and Management of CKD recommended considering "planning for preemptive kidney transplantation or dialysis access in adults when the GFR is < 15-20ml/min per 1.734 m² or risk of kidney replacement therapy (KRT) is greater than 40% over 2 years."

REGULATORY STATUS

Kidney transplants, as surgical procedures, do not require U.S. Food and Drug Administration (FDA) approval.

The Health Resources Services Administration (HRSA) oversees the transplantation of vascularized human organs through the Organ Procurement and Transplantation Network (OPTN), which was established through the National Organ Transplant Act (NOTA) approved by U.S. Congress in October 1984. It served as a model for development of other transplant networks worldwide, directing how organ allocation would be managed on a national basis. The model was developed through a unique public-private partnership with the United Network for Organ Sharing (UNOS). For more information view the UNOS website, available from: <https://unos.org/about/> [accessed 2026 Feb 26].

Medical Policy: Kidney Transplantation

Policy Number: 07.02.04

Page: 7 of 10

The HIV Organ Policy Equity (HOPE) Act was enacted in November 2013 and amended regulations of NOTA. The amendment removed clinical research and institutional review board (IRB) requirements for transplantation of kidneys and livers from donors with HIV to recipients with HIV.

CODE(S)

- Codes may not be covered under all circumstances.
- Code list may not be all inclusive (AMA and CMS code updates may occur more frequently than policy updates).
- (E/I)=Experimental/Investigational
- (NMN)=Not medically necessary/appropriate

CPT Codes

| Code | Description |
|-------|--|
| 50300 | Donor nephrectomy (including cold preservation); from cadaver donor, unilateral or bilateral |
| 50320 | Donor nephrectomy (including cold preservation); open, from living donor |
| 50323 | Backbench standard preparation of cadaver donor renal allograft prior to transplantation, including dissection and removal of perinephric fat, diaphragmatic and retroperitoneal attachments, excision of adrenal gland, and preparation of ureter(s), renal vein(s), and renal artery(s), ligating branches, as necessary |
| 50325 | Backbench standard preparation of living donor renal allograft (open or laparoscopic) prior to transplantation, including dissection and removal of perinephric fat and preparation of ureter(s), renal vein(s), and renal artery(s), ligating branches, as necessary |
| 50327 | Backbench reconstruction of cadaver or living donor renal allograft prior to transplantation; venous anastomosis, each |
| 50340 | Recipient nephrectomy (separate procedure) |
| 50360 | Renal allotransplantation, implantation of graft; without recipient nephrectomy |
| 50365 | Renal allotransplantation, implantation of graft; with recipient nephrectomy |
| 50370 | Removal of transplanted renal allograft |
| 50380 | Renal auto-transplantation, reimplantation of kidney |

Copyright © 2026 American Medical Association, Chicago, IL

Medical Policy: Kidney Transplantation

Policy Number: 07.02.04

Page: 8 of 10

HCPCS Codes

| Code | Description |
|----------------|-------------|
| Not Applicable | |

ICD10 Codes

| Code | Description |
|-------------|---|
| N18.1-N18.9 | Chronic kidney disease (CKD) (code range) |

REFERENCES

- American Society of Transplant Surgeons (ASTS), The American Society of Transplantation (AST), The Association of Organ Procurement Organizations (AOPO) [Internet] Statement on transplantation of organs from HIV-infected deceased donors.[accessed 2026 Feb 25] Available from: https://asts.org/docs/default-source/position-statements/transplantation-of-organs-from-hiv-infected-deceased-donors-july-22-2011.pdf?sfvrsn=fbae5a20_4
- Black CK, et al. Solid organ transplantation in the 21st century. *Ann Transl Med.* 2018 Oct;6(20):409.
- Chaudhry D, et al. Survival for waitlisted kidney failure patients receiving transplantation versus remaining on waiting list: systematic review and meta-analysis. *BMJ.* 2022 Mar 01;376:e068769.
- Choi A, et al. First porcine to human kidney transplantation performed in the United States. *Artif Organs.* 2022 Feb;46(2):178-179.
- Delgado C, et al. A unifying approach for GFR estimation: Recommendations of the NKF-ASN task force on reassessing the inclusion of race in diagnosing kidney disease. *Am J Kidney Dis.* 2022 Feb;79(2):268-288.e1.
- Gupta M, et al. Repeat kidney transplantation after failed first transplant in childhood: past performance informs future performance. *Transplantation.* 2015 Aug;99(8):1700-8.
- Kainz A, et al. Waiting time for second kidney transplantation and mortality. *Clin J Am Soc Nephrol.* 2022 Jan;17(1):90-97.
- Kute VB, et al, Outcomes of living donor kidney transplant after sars-cov-2 infection in both the donor and the recipient: a multicenter study. *Exp Clin Transplant.* 2022 Oct;20(10):908-916.
- Lentine KL, et al. OPTN/SRTR 2023 annual data report: kidney. *Am J Transplant.* 2025 Feb;25(2):S22-S137.
- Meshram KV, et al. Update on coronavirus 2019 vaccine guidelines for transplant recipients. *Transplant Proc.* 2022 Jul-Aug;54(6):1399-1404.
- National Kidney Foundation. [Internet] Glomerular filtration rate (GFR). [updated 2026 Feb 05;

Medical Policy: Kidney Transplantation

Policy Number: 07.02.04

Page: 9 of 10

accessed 2026 Feb 25]. Available from: <https://www.kidney.org/atoz/content/gfr>

Navaneethan SD, et al. KDOQI US commentary on the KDIGO 2024 clinical practice guideline for the evaluation and management of CKD. Am J Kidney Dis. 2025 Feb;85(2):135-176.

Organ Procurement and Transplant Network (OPTN). [Internet]. [accessed 2026 Feb 26]. Available from: <https://www.hrsa.gov/optn?from=optn.transplant.hrsa.gov>

Philipse DM, et al. Timing of the pre-transplant workup for renal transplantation: is there room for improvement? Clin Kidney J. 2022 Jan 17;15(6):1100-1108.

Porrett PM, et al. First clinical-grade porcine kidney xenotransplant using a human decedent model. Am J Transplant 2022 Jan 20.

Saeed B. Kidney retransplantation in children. Exp Clin Transplant. 2024 Jan;22 (Suppl 1):37-43.

Steinman TI, et al. The clinical practice committee of the American Society of Transplantation. Guidelines for the referral and management of patients eligible for solid organ transplantation. Transplant 2001;71(9):1189-204.

Trullas JC, et al. Renal transplantation in HIV-infected patients: 2010 update. Kidney Int. 2011 Apr;79(8):825-42.

United Network for Organ Sharing (UNOS) [Internet]. How we match organs. 2026 Jan 12. [accessed 2026 Feb 25]. Available from: <https://unos.org/transplant/how-we-match-organs/>

United Network for Organ Sharing (UNOS) [Internet]. [accessed 2026 Feb 26]. Available from: <https://unos.org/about/>

Wallisch SS, et al. Survival benefit of first single-organ deceased donor kidney transplantation compared with long-term dialysis across ages in transplant-eligible patients with kidney failure. JAMA Netw Open. 2022 Oct 3;5(10):e2234971.

SEARCH TERMS

Not Applicable

CENTERS FOR MEDICARE AND MEDICAID SERVICES (CMS)

There is currently no National Coverage Determination (NCD) or Local Coverage Determination (LCD) for Kidney Transplantation.

Please refer to the following Medicare Benefit Policy Manual Chapter 11-End Stage Renal Disease (ESRD). Revised 257, 03/01/19. [Medicare Benefit Policy Manual](#) [accessed 2026 Feb 26]

PRODUCT DISCLAIMER

- Services are contract dependent; if a product does not cover a service, medical policy criteria do not apply.
- If a commercial product (including an Essential Plan or Child Health Plus product) covers a specific service, medical policy criteria apply to the benefit.

Medical Policy: Kidney Transplantation

Policy Number: 07.02.04

Page: 10 of 10

- If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit.
- If a Medicare product (including Medicare HMO-Dual Special Needs Program (DSNP) product) covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.
- If a Medicare HMO-Dual Special Needs Program (DSNP) product DOES NOT cover a specific service, please refer to the Medicaid Product coverage line.

| POLICY HISTORY/REVISION | |
|--|--|
| Committee Approval Dates | |
| 04/19/00, 08/16/01, 06/20/02, 05/21/03, 02/19/04, 02/17/05, 01/19/06, 02/15/07, 01/17/08, 03/19/09, 03/18/10, 03/17/11, 02/16/12, 01/17/13, 02/20/14, 02/19/15, 03/17/16, 03/16/17, 03/15/18, 03/21/19, 03/19/20, 03/18/21, 03/17/22, 05/18/23, 04/18/24, 04/17/25, 04/16/26 | |
| Date | Summary of Changes |
| 04/16/26 | <ul style="list-style-type: none">• Annual review; policy intent unchanged. |
| 04/17/25 | <ul style="list-style-type: none">• Annual review; policy intent unchanged |
| 01/01/25 | <ul style="list-style-type: none">• Summary of changes tracking implemented. |
| 04/19/00 | <ul style="list-style-type: none">• Original effective date |