

MEDICAL POLICY

Medical Policy Title	Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members
Policy Number	7.01.121
Current Effective Date	January 1, 2027
Next Review Date	September 2027

Our medical policies are guides to evaluate technologies or services for medical necessity. Criteria are established through the assessment of evidence based, peer-reviewed scientific literature, and national professional guidelines. Federal and state law(s), regulatory mandates and the member's subscriber contract language are considered first in the determination of a covered service.

(Link to [Product Disclaimer](#))

SECTION LINKS

Hip Arthroplasty

[Partial](#)

[Total](#)

[Revision](#)

[Resurfacing](#)

[Osteotomy-Pediatrics](#)

Hip Arthroscopy

[Hip Arthroscopy](#)

[Hip Labral Repair](#)

POLICY STATEMENT(S)

Hip Arthroplasty

Partial Hip Arthroplasty

- I. Partial hip replacement is medically necessary when ALL the following are met:
 - A. Body Mass Index (BMI) is either:
 1. 40 kg/m² or less, or
 2. If BMI is greater than 40 kg/m², documented weight reduction of at least 10% before surgery;
 - B. ANY of the following conditions:
 1. Comminuted femoral head or femoral neck fracture
 2. Completely displaced femoral neck or femoral head fracture;

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 2 of 32

3. Impacted femoral neck or femoral head fracture;
4. Partially displaced femoral neck or femoral head fracture;
5. Intracapsular fracture that is non-displaced for which fixation is not considered an appropriate treatment option;

C. Traumatic hip injury.

Total Hip Arthroplasty- Outpatient

- II. Outpatient total hip arthroplasty is medically necessary when BOTH the following are met:
- A. BMI of 40 kg/m² or less; and
 - B. Presence of ANY of the following conditions when associated criteria are met:
 1. Developmental dysplasia of the hip;
 2. Displaced or comminuted femoral neck or femoral head;
 3. Post-traumatic arthritis;
 4. Malunion/non-union femoral head or proximal femur fracture;
 5. Prior fracture treated with open reduction internal fixation (ORIF) resulting in advanced joint disease;
 6. Osteonecrosis (Avascular Necrosis [AVN]) with subchondral fracture or crescent sign demonstrated on imaging;
 7. Symptomatic Osteonecrosis/AVN with stage IV collapse; or
 8. Osteoarthritis with ALL the following:
 - a. Arthritis type is any of the following:
 - i. Arthritis is Tonnis grade 2 or 3;
 - ii. Rheumatoid arthritis;
 - iii. Post-traumatic arthritis;
 - iv. Inflammatory arthropathy; and
 - b. Hip pain causing loss of function for 12 weeks or more that interferes with activities of daily living; and
 - c. Failure of conservative therapy to improve symptoms or functional status and includes BOTH of the following:
 - i. A physical therapy program (including telehealth-based rehabilitation when appropriate) that is provided by a physical therapist or chiropractor for at least 6 weeks in the past six 6 months; and

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 3 of 32

- ii. ANY of the following treatments:
 - a) Modification of activity/protective weight-bearing/use of an orthotic;
 - b) Intra-articular corticosteroid injection;
 - c) Medication management; and
- d. Diagnostic imaging studies (e.g., radiographs, CT, and/or MRI as clinically indicated) demonstrate ANY of the following findings:
 - i. Bone-on-bone articulation;
 - ii. Severe joint-space narrowing;
 - iii. Advanced osteoarthritis;
 - iv. Tönnis Grade IV deformity;
 - v. Symptomatic Osteonecrosis/AVN with stage IV collapse; and
- e. Documentation demonstrates ANY of the following symptoms:
 - i. Contracture;
 - ii. Antalgic gait or Trendelenburg gait;
 - iii. Pain with hip rotation;
 - iv. Leg-length discrepancy;
 - v. Reduction in range of motion.

Total Hip Arthroplasty- Inpatient

- III. An inpatient total hip arthroplasty is medically necessary when ALL the following are met:
- A. Criteria for outpatient total hip arthroplasty are met (P.S. II); and
 - B. Presence of ANY of the following perioperative risk factors:

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 4 of 32

1. Congestive heart failure;
2. Chronic atrial fibrillation that is not adequately controlled;
3. Transient ischemic attack (TIA) or stroke within the preceding three months;
4. Diabetes mellitus with HbA1c greater than 8% within the last 90 days;
5. Stage 4 or 5 chronic kidney disease;
6. End-stage renal disease;
7. Dialysis dependence;
8. eGFR less than 30 mL/min/1.73 m² within 30 days; or
9. Serum creatinine greater than 3 mg/dL within 30 days;
10. Inherited or acquired coagulation disorders;
11. Platelet count below 100,000 within the last 30 days;
12. Clinically significant abnormalities in coagulation testing;
13. Ongoing anticoagulation without reversal; or
14. Significant history of bleeding complications;
15. Significant chronic obstructive pulmonary disease (COPD) or emphysema; or
16. Home oxygen dependence;
17. Dementia or other cognitive impairment;
18. Schizophrenia;
19. Bipolar disorder;
20. Recurrent major depressive disorder; or
21. Other conditions that may increase postoperative risk or impair recovery;
22. Age 85 years or older;
23. Proximal femoral or adjacent soft tissue malignancy;
24. History of alcohol withdrawal seizures or delirium tremens;
25. Evidence of active drug toxicity manifested by physiologic instability, such as tachycardia, hypertension, or electrocardiographic abnormalities; or
26. American Society of Anesthesiologists (ASA) Physical Status Classification greater than 3.

IV. Total hip arthroplasty is not medically necessary for ANY of the following contraindications:

- A. Diabetes with HbA1c greater than 8%;

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 5 of 32

- B. Nicotine or tobacco use, including cigars, cigarette, e-cigarette, nicotine pouch, smokeless tobacco, or vape pen, within six (6) weeks of surgery;
- C. Intra-articular corticosteroid injection within 12 weeks of surgery;
- D. Clinically unstable or unmanaged comorbid conditions;
- E. Active infection, in treatment or is ongoing;
- F. Uncontrolled hypertension;
- G. Allergy to medical treatment material;
- H. Drug dependency that is not being appropriately managed.

Revision Hip Arthroplasty

- V. Revision hip arthroplasty is medically necessary when ALL the following are met:
 - A. BMI is either:
 - 1. 40 kg/m² or less, or
 - 2. If BMI is greater than 40 kg/m², documented weight reduction of at least 10% before surgery;
 - B. Prosthesis is painful, unstable, or causing loss of function;
 - C. ANY of the following:
 - 1. Recurrent instability, dislocation, or subluxation;
 - 2. Intolerable leg-length discrepancy;
 - 3. Aseptic loosening;
 - 4. Osteolysis;
 - 5. Implant component failure;
 - 6. Implant recall;
 - 7. Symptomatic synovitis or metallosis;
 - 8. Prosthetic infection or failure requiring removal.

Hip Resurfacing

- VI. Hip resurfacing is medically appropriate, when ALL the following are met:
 - A. The individual is age 55 years or less;
 - B. Active lifestyle;
 - C. Candidate for conventional total hip arthroplasty;
 - D. Femoral head size of greater than or equal to 46mm;

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 6 of 32

- E. Adequate femoral bone stock;
 - F. Presurgical x-rays within 2 to six 6 months;
 - G. Failure of conservative therapy including ALL the following:
 - 1. A physical therapy program (including telehealth- based rehabilitation when appropriate) that is provided by a physical therapist or chiropractor for at least 6 weeks in the past 6 months; and
 - 2. NSAID therapy for a minimum of 4 weeks or attempts with non-medication interventions such as limitation of weight-bearing activity, application of rest, ice and heat, splints, and exercise.
- VII. Hip resurfacing is not medically necessary for ANY of the following contraindications:
- A. Developmental abnormalities affecting the acetabulum or femur;
 - B. Systemic infection or other active infectious process;
 - C. Immunocompromised status, including chronic corticosteroid therapy;
 - D. Clinically significant leg length inequality;
 - E. Known hypersensitivity or allergy to implant metal components;
 - F. Osteonecrosis/AVN affecting more than 50% of the femoral head;
 - G. Pregnancy;
 - H. Moderate-to-severe renal dysfunction, including an estimated glomerular filtration rate (eGFR) below 60 mL/min/1.73m²;
 - I. Severe reduction in bone quality or density, such as advanced osteopenia or osteoporosis;
 - J. History of slipped capital femoral epiphysis (SCFE); or
 - K. Incomplete skeletal maturity.

Osteotomy- Pediatrics

- VIII. Pelvic and/or femoral osteotomy for the treatment of developmental dysplasia of the hip in a pediatric individual is medically necessary when documentation supports ALL the following:
- A. Preoperative imaging performed within the previous 4 weeks demonstrates findings consistent with developmental dysplasia of the hip and includes either of the following:
 - 1. Hip arthrography; or
 - 2. MRI, or CT when MRI is contraindicated;
 - B. At least two (2) of the following clinical features are present:
 - 1. Positive Barlow or Ortolani examination findings;

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 7 of 32

2. Positive Galeazzi sign;
 3. Asymmetric gluteal and/or thigh skin folds;
 4. Limb-length discrepancy
 5. Limited or asymmetric hip abduction with associated mild hip instability; or
 6. Limping or waddling gait in an ambulatory child;
- C. Failure of ANY of the following age-appropriate conservative treatment approaches:
1. Closed reduction with spica cast immobilization for 12 to 18 weeks;
 2. Open reduction followed by spica cast immobilization for 6 to 12 weeks; or
 3. Pavlik harness treatment for 4 to 12 weeks in infants younger than 6 months of age.

Hip Arthroscopy

Hip Arthroscopy- Diagnostic

IX. Diagnostic hip arthroscopy is infrequently indicated, and therefore is only medically necessary when ALL the following are met:

- A. Hip pain for at least 6 months that results in functional impairment that interferes with activities of daily living;
- B. Conventional imaging has not established a diagnosis, including both MRI and plain radiographs; and
- C. Documentation demonstrates failure of conservative therapy including both of the following:
 1. A supervised course of physical therapy, including telehealth-based rehabilitation when appropriate, completed for a minimum of six (6) weeks within the preceding six (6) months; and
 2. ANY of the following:
 - a. Activity modification, protected weight-bearing, or orthotic support;
 - b. Intra-articular corticosteroid injection; or
 - c. Pharmacologic pain management (e.g., nonsteroidal anti-inflammatory drugs [NSAIDs] or analgesics).

Hip Arthroscopy-Biopsy, Foreign Body Removal, Synovectomy

X. Hip arthroscopy biopsy, foreign body removal, synovectomy, or a combination of these procedures is medically necessary when ALL the following are met:

- A. Hip pain for at least 3 months;
- B. Both of the following symptoms:

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 8 of 32

1. Pain results in functional impairment that interferes with activities of daily living; and
 2. Is associated with mechanical symptoms such as catching, clicking, grinding, locking, or popping;
- C. Diagnostic imaging studies (e.g., radiographs, CT, and/or MRI as clinically indicated) demonstrate ANY of the following:
1. Calcification;
 2. Inflammatory changes;
 3. Intra-articular loose body;
 4. Nodular lesion(s);
 5. Pannus formation;
 6. Synovial proliferation;
- D. Clinical exam reveals restricted range of motion of the affected hip;
- E. Failure of conservative therapy including ALL the following:
1. A physical therapy program (including telehealth-based rehabilitation when appropriate) that is provided by a physical therapist or chiropractor for at least 6 weeks in the past 6 months, and
 2. ANY of the following treatment modalities:
 - a. Activity modification, protected weight-bearing, or orthotic use;
 - b. Intra-articular corticosteroid injection; or
 - c. Pharmacologic pain management.

Hip Arthroscopy- Femoroacetabular Impingement

XI. Arthroscopic treatment of CAM-type or Pincer-type femoroacetabular impingement is medically necessary when ALL the following criteria are met:

- A. BMI is 40kg/m² or less;
- B. Imaging confirms both the following:
 1. The presence of either of the following:
 - a. CAM-type impingement, demonstrated by femoral head-neck junction abnormalities (e.g., loss of femoral head sphericity, pistol-grip deformity, or alpha angle greater than 55 degrees); or
 - b. Pincer-type impingement, demonstrated by acetabular overcoverage findings such as acetabular retroversion, crossover sign, acetabular protrusion, posterior wall sign, prominent ischial spine sign, or center-edge angle greater than 35 degrees;

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 9 of 32

and

2. The absence of either of the following on weight-bearing radiographs:
 - a. Moderate-to-severe osteoarthritis (Tönnis grade 2 or 3); or
 - b. Hip dysplasia; and
- C. Physical exam demonstrates the following:
 1. Hip and groin pain consistent with impingement, including pain with flexion, adduction, and internal rotation; and
 2. Symptoms are reproducible or exacerbated by hip positioning;
Mechanical symptoms such as locking, catching, grinding, or popping and positive provocative maneuvers such as FABER or Fitzgerald testing may also be present;
and
- D. Failure of conservative therapy including ALL the following:
 1. A physical therapy program (including telehealth-based rehabilitation when appropriate) that is provided by a physical therapist or chiropractor for at least 6 weeks in the past six 6 months, and
 2. ANY of the following treatment modalities:
 - a. Activity modification, protected weight-bearing, or orthotic use;
 - b. Intra-articular corticosteroid injection; or
 - c. Pharmacologic pain management.

Hip Labral Repair

XII. Hip labral repair is medically necessary when ALL the following are met:

- A. Imaging findings include both the following:
 1. Weight-bearing radiographs do not demonstrate moderate-to-severe osteoarthritis (Tönnis grade 2 or 3) or hip dysplasia;
 2. Magnetic resonance imaging confirms an acetabular labral tear;
- B. Hip pain for at least 6 weeks that includes all the following:
 1. The pain results in functional impairment that interferes with activities of daily living;
 2. The pain is accompanied by examination findings consistent with impingement, including pain with flexion, adduction, and internal rotation.
Mechanical symptoms such as locking, catching, grinding, or popping may also be present.

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 10 of 32

- C. Documentation requirements for conservative treatments are EITHER of the following:
 - 1. Documentation is not required for the following conditions:
 - a. Irreparable or substantially deficient labral tissue;
 - b. A labral tear identified during another medically necessary hip arthroscopic procedure;
 - c. Loose bodies, foreign bodies, or synovial pathology requiring surgical removal;
 - d. Advanced structural femoroacetabular impingement with clearly associated labral pathology;
 - e. Suspected or confirmed septic arthritis;
 - OR
- D. Documentation demonstrates failure of conservative therapy following:
 - 1. A supervised physical therapy program, including telehealth-based rehabilitation when appropriate, for a minimum of 6 weeks within the preceding 6 months; and
 - 2. ANY of the following:
 - a. Activity modification, protected weight-bearing, or orthotic support;
 - b. Intra-articular corticosteroid injection; or
 - c. Pharmacologic pain management.

Not Medically Necessary Indications

XIII. The following indications are Not Medically Necessary:

- A. Hip Arthroscopy for ANY of the following conditions:
 - 1. Bone marrow edema;
 - 2. Radiographic or clinical evidence of fracture;
 - 3. Intra-articular loose body or synovial pathology (see procedure specific criteria for arthroscopic removal of loose bodies and synovial pathology in Policy Statement XI);
 - 4. Tear involving the abductor, adductor, or hamstring tendon complex;
- B. Diagnostic Hip Arthroscopy when ANY of the following are present:
 - 1. Advanced degenerative joint disease when documentation demonstrates significant arthritic changes of the affected hip, including either of the following:
 - a. Joint space narrowing to 2 mm or less on radiographs; or
 - b. Subchondral edema identified on MRI;

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 11 of 32

2. Structural hip abnormalities with an established diagnosis and imaging confirms ANY of the following:
 - a. Acetabular protrusion;
 - b. Acetabular overcoverage involving the anterolateral rim;
 - c. Femoroacetabular impingement (FAI);
 - d. Alpha angle greater than 50 degrees;
 - e. Crossover sign;
 - f. Posterior wall sign;
 - g. Prominent ischial spine sign;
 - h. Abnormal femoral head contour resulting in loss of normal sphericity;
 - i. Prominent femoral head-neck junction (e.g., pistol-grip deformity); or
 - j. Acetabular retroversion with a center-edge angle greater than 35 degrees;
- C. Hip dysplasia when imaging findings are consistent, including ANY of the following, when the proposed procedure does not include a periacetabular osteotomy:
 1. Lateral center-edge angle less than 20 degrees;
 2. Anterior center-edge angle less than 20 degrees;
 3. Tönnis angle greater than 15 degrees; or
 4. Femoral head extrusion index greater than 25%;
- D. Alternative identifiable pathology that adequately explains symptoms, including ANY of the following:
 1. Intra-articular loose body;
 2. Pigmented villonodular synovitis (PVNS);
 3. Synovial chondromatosis;
 4. Ischiofemoral impingement;
 5. Pubic edema or osteitis pubis; or
 6. MRI-confirmed tear involving the labrum, adductor tendon, hamstring tendon, gluteus medius, or gluteus minimus.
- E. Presence of active infection; or
- F. Unmanaged substance use disorder/dependency.

RELATED POLICIES

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 12 of 32

Corporate Medical Policy

Not Applicable

POLICY GUIDELINE(S)

Not Applicable

DESCRIPTION

Total Hip Arthroplasty

Total hip replacement is a surgical technique in which the femoral head and neck are removed, and the femoral canal (marrow space) is reamed out. The damaged hip joint is replaced with an artificial prosthesis composed of two or three different components: (1) the head that replaces the original femoral head; (2) the femoral component (a metal stem placed into the femur); and (3) the acetabular component, which is implanted into the acetabulum. The stem may be secured using bone cement or press-fit for the bone to grow into it.

The Tonnis Classification System is commonly used to describe the presence of osteoarthritis in the hips, with grading as follows:

- Grade 0: No signs of osteoarthritis.
- Grade 1: Sclerosis of the joint, with slight joint space narrowing and osteophyte formation and no or slight loss of femoral head sphericity.
- Grade 2: Small cysts in the femoral head or acetabulum, with moderate joint space narrowing and moderate loss of femoral head sphericity.
- Grade 3: Large cysts in the femoral head or acetabulum, severe joint space narrowing or obliteration of the joint space, and severe deformity and loss of sphericity of the femoral head.

Revision of hip replacement (partial or total) involves surgical reconstruction or replacement due to failure or complications of previous hip replacement.

Non-surgical management regarding the treatment of hip osteoarthritis is defined as any provider-directed, non-surgical treatment that has been demonstrated in the scientific literature to be efficacious and/or is considered reasonable care in the treatment of hip pain from osteoarthritis. The types of treatment can include but are not limited to relative rest/activity modification, weight loss, supervised physiotherapy modalities and therapeutic exercises, oral prescription and non-prescription medications, assistive devices (e.g., cane, crutches, walker, wheelchair), and/or intra-articular (i.e., steroid) injections.

Hip Resurfacing

Total hip resurfacing is an alternative to watchful waiting or total hip arthroplasty (THA) for younger, active individuals less than 65 years old with hip disease such as osteoarthritis, rheumatoid arthritis, or advanced avascular necrosis.

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 13 of 32

In total hip resurfacing, the surface of the femoral head is trimmed and covered with a hollow metal hemisphere that fits into a metal acetabular cup. It is believed to optimize stress transfer to the proximal femur, and, because of the large diameter of the articulation, to offer stability and optimal range of movement. Because total hip resurfacing preserves the proximal femoral bone stock, it may not compromise future total hip replacements.

Proposed advantages of total hip resurfacing as compared with THA include preservation of the femoral neck and femoral canal, thus allowing for revision or conversion to THA, if required. In addition, the resurfaced head is more similar in size to the normal femoral head, thus increasing the stability and decreasing the risk of dislocation compared with THA.

Non-surgical management with regard to the treatment of lower extremity joint pain is defined as any provider-directed, non-surgical treatment which has been demonstrated in scientific literature to be efficacious and/or is considered reasonable care in the treatment of lower extremity joint pain. Types of treatment may include but are not limited to relative rest/activity modification, weight loss, supervised physiotherapy modalities and therapeutic exercises, oral prescription and non-prescription medications, assistive devices (e.g., cane, crutches, walker, wheelchair), and/or intra-articular injections (e.g., steroid).

Total hip resurfacing using polyethylene components offers a promising bone-conserving alternative to total hip replacement, particularly for individuals with metal allergies or small femoral geometries (variations in the shape and size of the upper part of the femur, particularly around the hip joint). Earlier concerns with conventional polyethylene relative to wear, thickness, osteolysis, and deformation are currently being studied with the use of cross-linked polyethylene which provides a low-friction bearing surface, mimicking the natural joint's articulation and thus promoting good outcomes.

SUPPORTIVE LITERATURE

Total Hip Arthroplasty

In a meta-analysis, Smith and colleagues (2010) compared the clinical and radiological outcomes and complication rates of hip resurfacing (HRS) and total hip arthroplasty (THA). A systematic review was undertaken of all published (Medline, CINAHL, AMED, EMBASE) and unpublished or gray literature research databases up to January 2010. Clinical and radiological outcomes, as well as complications of HRS, were compared to those of THA, using risk ratio, mean difference, and standardized mean difference statistics. Studies were critically appraised using the CASP appraisal tool. A total of 46 studies were identified from 1,124 citations. These included 3,799 HRSs and 3,282 THAs. On meta-analysis, functional outcomes for subjects following HRS were better than or the same as for subjects with a THA, but there were statistically significant increases in incidence of heterotopic ossification, aseptic loosening, and revision surgery with HRS, compared to THA. The evidence base showed a number of methodological inadequacies, such as the limited use of power calculations and poor or absent blinding of both patients and assessors, possibly giving rise to assessor bias. The authors concluded that, on the basis of the current evidence base, HRS may have better functional outcomes

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 14 of 32

than THA, but the increased risks of heterotopic ossification, aseptic loosening, and revision surgery following HRS indicate that THA is superior in terms of implant survival.

In a 2019 retrospective cohort study, Inoue et al compared post-operative complications and survivorship of total hip and knee arthroplasty in dialysis and renal transplantation patients. They included a total of 107 patients undergoing primary total joint arthroplasty (TJA), including 50 who were receiving dialysis and 57 who had a prior renal transplantation. The end point was defined as revision surgery secondary to post-operative complications. Researchers found a significantly higher rate of post-operative complications in the dialysis cohort (28%, n=14 of 50 joints) compared to the renal transplant cohort (7.1%, n= 4 of 57 joints). There was a higher rate of surgical site infection (SSI) and periprosthetic joint infection (PJI) in dialysis patients, compared with renal transplantation patients (18% versus 3.5%, P=0.02). In addition, there was an increased rate of revision surgery in the dialysis cohort, compared to transplant cohort (24% versus 3.5%, P=0.002). A multi-variate analysis considering demographics and comorbidities revealed that patients with renal transplantation were less likely to have revision surgery, compared to patients on dialysis as the time of arthroplasty (95 % CI, P=0.031), and demonstrated a strong trend for lower complications (95% CI, P=0.76), although the latter was not statistically significant. Researchers concluded that transplantation was independently associated with reduced rates of revision surgery in the setting of chronic renal failure, suggesting that those who are candidates may benefit from renal transplantation before undergoing elective TJA.

Chou et al (2021) compared outcomes of TJA, including THA and total knee arthroplasty, in individuals with end-stage renal disease who were either receiving dialysis or had undergone a kidney transplant (KT). The analysis included 22 studies encompassing 9,384 patients (n = 8,921 dialysis; n = 463 KT). The overall mortality rate after TJA was 14.9%-13.8% in dialysis patients and 15.8% in KT patients. The overall surgical site complication (SSC) rate was 3.3%, with KT patients experiencing a rate of 3.6%. SSCs included hematoma, seroma, delayed wound healing, and superficial infections. The overall PJI rate was 3.9%, with dialysis patients at 4.0% and KT patients at 3.7%. Multivariate regression analysis found no statistically significant differences in outcomes between dialysis and KT patients. Both groups demonstrated elevated risks compared to the general population, with comparable outcomes between them.

Kim et al (2022) conducted a systematic review and meta-analysis of 10 retrospective studies that included 3,631,861 patients and 14,996 solid organ transplantations (SOT). They evaluate whether patients with a history of SOT experienced different outcomes after primary hip arthroplasty compared to non- SOT patients. Medical complications included cardiac complications, pneumonia, acute kidney injury, and pulmonary thromboembolism and deep vein thrombosis. The study showed that patients with a history of SOT undergoing hip arthroplasty are at higher risk for medical complications, transfusion, readmission, and short-term mortality compared to non-transplant patients. However, surgical complication rates such as infection and revision are not significantly different.

Hip Resurfacing

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 15 of 32

Evidence from numerous case series demonstrates symptomatic and functional improvements that appear to be comparable to those obtained with the current generation of total hip replacement in individuals younger than age 65 years at similar follow-up duration. In addition, hip resurfacing leaves femoral bone stock intact and therefore revision is technically similar to primary total hip replacement. Increased concentrations of metal ions have been documented after metal-on-metal hip resurfacing in some individuals. The effect of metal ion release on a fetus is also unknown.

Za and colleagues (2024) performed a systematic review and meta-analysis of eight (8) randomized control trials that compared the outcomes of hip resurfacing (HR) and total hip arthroplasty (THA) in the treatment of hip osteoarthritis in 844 patients (n=387 hip resurfacing; n=469 THA). The mean follow-up was 7.72 years. The systematic review identified no significant differences between the groups in terms of Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) and University of California Los Angeles (UCLA) scores, revision rates, infection, aseptic loosening, or pseudotumor occurrence (all $p > 0.05$). However, the dislocation rate was significantly lower in the hip resurfacing group (0.72% vs 4%; $p = 0.04$). Some studies evaluated metal ion levels with slightly higher chromium levels noted in THA patients postoperatively. The most common cause of revision in HR was aseptic loosening (48%), while adverse reactions to metal debris were the leading cause in THA (30.77%). The authors concluded that HR is a safe and effective alternative to THA.

In a 2019 retrospective cohort study, Inoue and colleagues compared post-operative complications and survivorship of total hip and knee arthroplasty in dialysis and renal transplantation patients. They included a total of 107 patients undergoing primary total joint arthroplasty, including 50 who were receiving dialysis and 57 who had a prior renal transplantation. The end point was defined as revision surgery secondary to post-operative complications. Researchers found a significantly higher rate of post-operative complications in the dialysis cohort (28%, n=14 of 50 joints), compared to the renal transplant cohort (7.1%, n= 4 of 57 joints). There was a higher rate of SSI and PJI in dialysis patients, compared with renal transplantation patients (18% versus 3.5%, $P = 0.02$). In addition, there was an increased rate of revision surgery in the dialysis cohort, compared to transplant cohort (24% versus 3.5%, $P = 0.002$). A multi-variate analysis considering demographics and comorbidities revealed that patients with renal transplantation were less likely to have revision surgery, compared to patients on dialysis as the time of arthroplasty (95 % CI, $P = 0.031$) and demonstrated a strong trend for lower complications (95% CI, $P = 0.76$), although the latter was not statistically significant. Researchers concluded that transplantation was independently associated with reduced rates of revision surgery in the setting of chronic renal failure, suggesting that those who are candidates may benefit from renal transplantation before undergoing elective total joint arthroplasty.

Ren and colleagues (2024) performed a retrospective review of 104 individuals who underwent 134 hip resurfacing arthroplasties with either the Birmingham Hip Resurfacing (BHR) (n=67) implant or the ReCap Magnum implant (n=67) at a single institution between 2006 and 2018. The primary aim of this study was to evaluate and compare early and midterm blood metal ion levels in hips implanted with BHR and ReCap devices as both contain cobalt-chromium (CoCr) metals. The study participants were matched 1:1 by sex, femoral head and acetabular cup sizes (± 2 millimeters), age at surgery (± 5 y), and year of surgery (± 5 y). The ReCap group had lower median metal ion levels compared to

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 16 of 32

the BHR group at 1-2 years (Co: 1.5 versus 1.9 parts per billion [ppb], $P=0.018$; Cr: 1.3 versus 2.8 ppb, $P=0.008$) and 3-5 years (Co: 1.1 versus 1.9 ppb, $P=0.001$; Cr: 1.2 versus 2.2 ppb, $P=0.003$) after surgery. Metal levels were identified in both study groups but potentially not clinically significant. Correlation analysis showed no significant associations between Co and Cr ion levels and pre- and postoperative patient-reported outcomes. Revisions for both study groups varied and included three (3) BHR hip revisions due to adverse reactions to metal debris as compared to two (2) ReCap hips revisions: one for instability and another for periprosthetic fracture. The authors concluded that the study identified that the BHR group had higher metal ion levels than the ReCap group at one to two (1-2) and three to five (3-5) years after surgery; however, these metal levels are still low and in line with prior studies.

Zuke and colleagues (2025) assessed the long-term outcomes of the Birmingham Hip Resurfacing (BHR) implant in a retrospective study of 224 male individuals under 60 years of age with osteoarthritis. The study followed the participants for an average of 14 years. Survivorship analysis revealed excellent long-term durability, with 96.0% of BHR implants remaining free from any revision and 97.4% free from aseptic revision at 15 years. Eight individuals required revision surgery, primarily due to infection ($n=3$), pseudotumor formation with elevated metal ions ($n=2$), and femoral component loosening ($n=2$). Median serum cobalt and chromium levels were 1.4 ppb and 1.5 ppb, respectively, with only two (2) cases of late-onset elevated metal ions requiring revision. Study participant-reported outcomes were similar between the cohort of individuals who received the BHR implant and the study participant group who received total hip arthroplasty (THA), with no significant differences in the modified Harris Hip Score (92.65 vs. 93.56; $p=0.44$) or long-term activity levels. The authors concluded that this cohort study demonstrated that metal on metal hip resurfacing arthroplasty using BHR implants is safe and effective with outcomes comparable to THA surgery.

PROFESSIONAL GUIDELINE(S)

Total Hip Arthroplasty

In the 2023 American Academy of Orthopaedic Surgeons adopted the evidence-based Clinical Practice Guidelines for the Management of Osteoarthritis of the Hip that state:

- “They strongly recommend conservative measures such as physical therapy, corticosteroid or hyaluronic acid injections, pharmacological management (NSAIDs).”

In the 2008 International Cartilage Repair Society Guidelines for the Management of Hip and Knee Osteoarthritis (OA), Part II, stated:

- “Patients with hip or knee OA who are not obtaining adequate pain relief and functional improvement from a combination of non-pharmacological and pharmacological treatment should be considered for joint replacement surgery. Replacement arthroplasties are effective, and cost-effective interventions for patients with significant symptoms, and/or functional limitations associated with a reduced health-related quality of life, despite conservative therapy.” (Zhang et al 2008)

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 17 of 32

Hip Resurfacing

Professional Society Name	Guideline/Version/Year	Summary of Content
Hip Society	2012 Algorithmic Approach to the Diagnosis and Management of Metal-on-Metal Arthroplasty	Awareness of adverse local tissue reactions to metal debris. All arthroplasty patients returning for follow-up should be queried for pain, discomfort, or compromise of function. Description of ideal candidate for metal on metal arthroplasty. Listing of contraindications for metal on metal arthroplasty.
National Institute for Health and Care Excellence (NICE)	2014 Updated Guidance on THA and Total Hip Resurfacing for End-stage Arthritis of the Hip	NICE advised that both THA and total hip resurfacing were options for treating end-stage arthritis of the hip. NICE concluded that THA was more effective and less costly than total hip resurfacing in all analyses, that the revision rate was the most important key driver of costs and quality-adjusted life years, and that because the predicted revision rate of THA was less than 5% at 10 years in the population for whom both THA and total hip resurfacing were suitable, the revision rate standard for total hip resurfacing should be the same as that for THA. NICE recommended specific

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 18 of 32

		prostheses for THA and total hip resurfacing only if the prostheses have revision rates of 5% or less at 10 years.
--	--	--

REGULATORY STATUS

Surgeries of the hip are procedures and, therefore, not regulated by the United States Food and Drug Administration (FDA). However, devices and instruments used during the surgery are regulated as medical devices. All resurfacing devices including related components require FDA approval before marketing and use in the United States to ensure they are safe and effective for human use. Refer to the FDA Medical Device website. Available from: <https://www.fda.gov/medical-devices> [accessed 2026 May 11]

The FDA lists the most serious type of medical device recalls as well as early alert communications about corrective actions being taken by companies that the FDA believes are likely to be the most serious type of recalls. Available from: [Medical Device Recalls | FDA](#) [accessed 2026 May 11]

The Birmingham Hip Resurfacing Device (BHR), a metal-on-metal system, received FDA premarket approval in May 2006. The Cormet Hip Resurfacing system, another metal-on-metal system, received FDA premarket approval in July 2007. Two additional metal-on-metal hip resurfacing systems have received approval: the Cormet Hip Resurfacing System (Corin) in 2007 and the Conserve Plus Total Hip Resurfacing System (MicroPort Orthopedics) in 2009. Both implants were approved for skeletally mature patients with either noninflammatory degenerative arthritis (e.g., osteoarthritis and avascular necrosis) or inflammatory arthritis (e.g., rheumatoid arthritis).

CODE(S)

- Codes may not be covered under all circumstances.
- Code list may not be all inclusive (AMA and CMS code updates may occur more frequently than policy updates).
- (E/I)=Experimental/Investigational
- (NMN)=Not medically necessary/appropriate

CPT Codes

Code	Description
27033	Arthrotomy, hip, including exploration or removal of loose or foreign body
27035	Denervation, hip joint, intrapelvic or extrapelvic intra-articular branches of sciatic, femoral, or obturator nerves

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 19 of 32

Code	Description
27050	Arthrotomy, with biopsy; sacroiliac joint
27052	Arthrotomy, with biopsy; hip joint
27054	Arthrotomy with synovectomy, hip joint
27062	Excision; trochanteric bursa or calcification
27090	Removal of hip prosthesis; (separate procedure)
27091	Removal of hip prosthesis; complicated, including total hip prosthesis, methylmethacrylate with or without insertion of spacer
27122	Acetabuloplasty; resection, femoral head (e.g., Girdlestone procedure)
27125	Hemiarthroplasty, hip, partial (e.g., femoral stem prosthesis, bipolar arthroplasty)
27130	Arthroplasty, acetabular and proximal femoral prosthetic replacement (total hip arthroplasty), with or without autograft or allograft
27132	Conversion of previous hip surgery to total hip arthroplasty, with or without autograft or allograft
27134	Revision of total hip arthroplasty; both components, with or without autograft or allograft
27137	Revision of total hip arthroplasty; acetabular component only, with or without autograft or allograft
27138	Revision of total hip arthroplasty; femoral component only, with or without allograft

Copyright © 2026 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
S2118	Metal-on-metal total hip resurfacing, including acetabular and femoral components

ICD10 Codes

Code	Description
M05.051- M08.959	Inflammatory polyarthropathies (hip) (code range)
M12.551 - M12.559	Traumatic arthropathy, hip (code range)

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 20 of 32

Code	Description
M16.0 - M16.9	Osteoarthritis of hip (code range)
M80.051A - M80.059S, M80.851A- M80.859S, M84.451A- M84.453S, M84.459A- M84.459S, M84.551A- M84.559S, M84.651A- M84.659S	Pathologic fracture of neck of femur (hip) (code range)
M84.750A- M84.759S	Atypical femoral fracture (code range)
M87.050	Idiopathic aseptic necrosis of pelvis
M87.051- M87.059, M87.150 M87.151- M87.159, M87.250 M87.251- M87.256, M87.350 M87.351- M87.353, M87.850 M87.851- M87.859, M90.551- M90.559	Osteonecrosis of femur, pelvis and thigh (code range)

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 21 of 32

Code	Description
M97.01XA-M97.02XS	Periprosthetic fracture around internal prosthetic hip joint (code range)
S72.001A-S72.26XS	Fracture of head and neck of femur (code range)

REFERENCES

Abdelaal, M.S., et al. Global Perspectives on Arthroplasty of Hip and Knee Joints. The Orthopedic Clinics of North America. 2020;51(2):169-176.

Adeyemo, A., et al. Implant Selection for Proximal Humerus Fractures The Orthopedic Clinics of North America. 2021; 52(2):167-175.

Adib, F., et al. Gluteal compartment syndrome: A systematic review and meta-analysis. Injury. 2021; 53(3):1209-1217.

Adrados M, et al. Institutional adherence to the American Association of Hip and Knee Surgeons body mass index guidelines lowers perioperative emergency department visits in primary total knee arthroplasty. J Arthroplasty. 2023 Jun;38(6S):S88-S93.

American Academy of Orthopaedic Surgeons (AAOS) [Internet]. Appropriate use criteria: Osteoarthritis of the hip: management. Dec 2017. [accessed 2026 May 11]. Available from: <https://www.orthoguidelines.org/go/auc/>

American Academy of Orthopaedic Surgeons (AAOS) [Internet]. Evidence-based clinical practice guideline: Management of osteoarthritis of the hip. Dec 2023. [accessed 2026 May 11]. Available from: <https://www.aaos.org/globalassets/quality-and-practice-resources/osteoarthritis-of-the-hip/oah-cpg.pdf>

Anzillotti, G., et al. Conservative vs. Surgical Management for Femoro-Acetabular Impingement: A Systematic Review of Clinical Evidence. Journal of Clinical Medicine. 2022;11(19):5852.

Ardon, A. E. Safety Considerations for Outpatient Arthroplasty. Clinics in Sports Medicine. 2022;41(2):281-289.

Azar, F. M., et al. 2021. Fractures of the Acetabulum and Pelvis. M. Beebee (Ed.). Campbell's Operative Orthopaedics (14), (pp. 2958-3030). Philadelphia, PA: Elsevier Inc.

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 22 of 32

Baal, J. D., et al. Imaging of Overuse Injuries of the Hip. Radiologic Clinics of North America. 2023; 61(2):191-201.

Bannuru RR, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. Osteoarthritis Cartilage. 2019 Nov;27(11):1578-1589.

Bartels S, et al. Total hip arthroplasty leads to better results after low-energy displaced femoral neck fracture in patients aged 55 to 70 years: a randomized controlled multicenter trial comparing internal fixation and total hip arthroplasty Journal of Bone Joint Surgery. 2022;Aug;104(15):1341-1351.

Bhandari M, et al. Total hip arthroplasty or hemiarthroplasty for hip fracture. N Engl J Med. 2019 Dec 5;381(23):2199-2208.

Biundo, J. J. (2020). Bursitis, Tendinitis, and Other Periarticular Disorders and Sports Medicine. L. Goldman & A. I. Schafer (Eds.). Goldman-Cecil Medicine (26), (pp. 1703-1709.e3). Philadelphia: Elsevier, Inc.

Blankstein M, et al. What factors increase revision surgery risk when treating displaced femoral neck fractures with arthroplasty: a secondary analysis of the HEALTH Trial. J Orthop Trauma. 2020 Nov;34 Suppl 3:S49-S54.

Bostrom, M. P. & Cross, M. B. (2020). The Neuromuscular Hip. D. J. Berry & J. R. Lieberman (Eds.). Surgery of the Hip (2), (pp. 965-977). Philadelphia: Elsevier, Inc.

Browne J, et al. Peritoneal dialysis does not carry the same risk as hemodialysis in patients undergoing hip or knee arthroplasty. J Bone Joint Surg. 2019;101(14):1271-77.

Busconi, B., et al. (2020). Hip Imaging. M. D. Miller & S. R. Thompson (Eds.). DeLee, Drez, & Miller's Orthopaedic Sports Medicine (5), (pp. 935-946. e2). Philadelphia: Elsevier, Inc.

Cantrell, C. K., et al. A review on management of insufficiency fractures of the pelvis and acetabulum the orthopedic clinics of North America. 2022;53(4): 431-443.

Carek, S.M. Hip and Knee Injuries. Clinics in Office Practice. 2020;47(1):115-131.

Carlo JO, et al. Perioperative implications of end-stage renal disease in orthopaedic surgery. J Am Acad Orthop Surg. 2015 Feb;23(2):107-18.

Cavanaugh PK, et al. Complications and mortality in chronic renal failure patients undergoing total joint arthroplasty: A comparison between dialysis and renal transplant patients. J Arthroplasty. 2016 Feb;31(2):465-72.

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 23 of 32

Chalmers BP, et al. Primary total hip arthroplasty for charcot arthropathy is associated with high complications but improved clinical outcomes. *J Arthroplasty*. 2018 Sept;33(9):2912-2918.

Chambers, H. F. & Sakoulas, G. (2020). *Staphylococcal Infections*. L. Goldman & A. I. Schafer (Eds.). *Goldman-Cecil Medicine*, (pp. 1862-1868.e2). Philadelphia: Elsevier, Inc.

Chammout G, et al. HOPE-Trial: hemiarthroplasty compared with total hip arthroplasty for displaced femoral neck fractures in octogenarians: a randomized controlled trial. *JB JS Open Access*. 2019 May 1;4(2):e0059.

Chang, C., Greenspan, A. & Gershwin, M.E.(2021). *Osteonecrosis*. G.S. Firestein & R.C. Budd (Eds.). *Firestein & Kelley's Textbook of Rheumatology* (11), (pp. 1852-1875.e7). Philadelphia, PA: Elsevier, Inc.

Chang, N., and King, G. J. Elbow Hemiarthroplasty for the Treatment of Distal Humerus Fractures. *The Orthopedic Clinics of North America*. 2020;51(2):265-277.

Chen W, et al. Direct anterior versus posterolateral approaches for clinical outcomes after total hip arthroplasty: a systematic review and meta-analysis. *J Orthop Surg Res*. 2020 Jun 23;15(1):231.

Cheney, C. W., et al. Radiofrequency ablation for chronic hip pain: a comprehensive, narrative review. *Pain Medicine*.2021; 22(S1):S14-S19

Chou TA, et al. Dialysis patients have comparable results to patients who have received kidney transplant after total joint arthroplasty: a systematic review and meta-analysis. *EFORT Open Rev*. 2021 Aug 10;6(8):618-628.

Cochran, H, et al. The Removal of Total Hip Arthroplasty From the Inpatient Only List Has Improved Patient Selection and Expanded Optimization Efforts. *J Arthroplasty*.2023 ;38:1-18.

Cohen DA, Montgomery SJ, Khoshbin A, et al. Treatment of geriatric acetabular fractures: a concise review of the literature. *Orthop Clin North Am*. 2021;52(4):323-333.

Crenshaw, A. H. (2021). *Tuberculosis and other Unusual Infections*. F. M. Azar & J. H. Beaty (Eds.). *Campbell's Operative Orthopaedics* (14), (pp. 868-888.e2). Philadelphia: Elsevier, Inc.

De Paula, F. J. A., Black, D. M. & Rosen, C. J. (2020). *Osteoporosis: Basic and Clinical Aspects*. S. Melmed & R. J. Auchus (Eds.). *Williams Textbook of Endocrinology* (14), (pp. 1256-1297.e10). Philadelphia: Elsevier, Inc.

Decker S, Winkelmann M, Knipp BS, et al. Pelvic ring fractures. In: Browner BD, Jupiter JB, Krettek C, Anderson PA, editors. *Skeletal trauma: basic science, management, and reconstruction*. 6th ed.

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 24 of 32

Philadelphia (PA): Elsevier; 2020. p. 1108-1195.

Domb BG, Bheem R, Shapira J, et al. Minimum five-year outcomes of hip resurfacing: propensity-score matched against total hip arthroplasty control groups. *J Arthroplasty*. 2021;36(6):2012-2015.

Douglas SJ, et al. Comparing primary total hip arthroplasty in renal transplant recipients to patients on dialysis for end-stage renal disease: a nationally matched analysis. *J Bone Joint Surgery Am*. 2021; 103:2215-20.

Dowsey MM, et al. Effect of bariatric surgery on risk of complications after total knee arthroplasty. *JAMA Network Open*. 2022 Apr;5(4):e226722.

Edelstein AI, Nassif NA, McDonald D, et al. Synovial diseases of the hip. In: Berry DJ, editor. *Surgery of the hip*. 2nd ed. Philadelphia (PA): Elsevier; 2020. p. 426-433.

Ekhtiari S, et al. Total hip arthroplasty versus hemiarthroplasty for displaced femoral neck fracture: a systematic review and meta-analysis of randomized controlled trials. *J Bone Joint Surg Am*. 2020 Sep 16;102(18):1638-1645.

Farrow L, Faulkner A, Reidy M, et al. Surgical management for developmental dysplasia of the hip. *Orthop Trauma*. 2020;34(6):324-331.

Ferreira GE, O'Keefe M, Zadro JR, et al. The effectiveness of hip arthroscopic surgery for the treatment of femoroacetabular impingement syndrome: a systematic review and meta-analysis. *J Sci Med Sport*. 2021;24(1):21-29.

Ford, M. C. (2021). Surface Replacement Hip Arthroplasty. F. M. Azar & J. H. Beaty (Eds.). *Campbell's Operative Orthopaedics* (14), (pp. 334-347.e2). Philadelphia: Elsevier, Inc.

Frisch NB, Jacobs JJ. Revision for metal-on-metal or taper corrosion failures. In: Berry DJ, editor. *Surgery of the hip*. 2nd ed. Philadelphia (PA): Elsevier; 2020. p. 1197-1211.

Fujiwara T, Ogura K, Healey J, et al. Periacetabular reconstruction following limb-salvage surgery for pelvic sarcomas. *J Bone Oncol*. 2021;31:100396. doi:10.1016/j.jbo.2021.100396.

Gillespie, P., Sabri, O., . . . Amr, R. (2022). Fragility fractures of the acetabulum. *Orthopaedics and Trauma*, 36(2), 111-119.

Goldblum JR, Folpe AL, Weiss SW. Miscellaneous benign soft tissue tumors and pseudotumors. In: Goldblum JR, Folpe AL, editors. *Enzinger and Weiss's soft tissue tumors*. 7th ed. Philadelphia (PA): Elsevier; 2020. p. 1085-1106.

Guyton JL. Hip pain in the young adult and hip preservation surgery. In: Azar FM, Beaty JH, editors. *Campbell's operative orthopaedics*. 14th ed. Philadelphia (PA): Elsevier; 2021. p. 356-404.e8.

Hankins DA, Korcek L, Richter DL, et al. Femoroacetabular impingement and management of labral

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 25 of 32

tears in the athlete. Clin Sports Med. 2021;40(2):259-270.

Harkness JW, Crockarell JR. Arthroplasty of the hip. In: Azar FM, Beaty JH, editors. Campbell's operative orthopaedics. 14th ed. Philadelphia (PA): Elsevier; 2021. p. 178-333.

Harris JD. Hip arthroscopy. In: Miller MD, Thompson SR, editors. DeLee, Drez, and Miller's orthopaedic sports medicine. 5th ed. Philadelphia (PA): Elsevier; 2020. p. 947-956.e3.

Harris JD, Lewis BD, Park KJ, et al. Hip dysplasia. Clin Sports Med. 2021;40(2):271-288.

Heck RK, Toy PC. General principles of tumors. In: Azar FM, Beaty JH, editors. Campbell's operative orthopaedics. 14th ed. Philadelphia (PA): Elsevier; 2021. p. 890-956.

Hermans MM, Droeghaag R, Curfs I, et al. Minimally invasive sacroiliac joint fusion versus conservative treatment in patients with sacroiliac joint dysfunction: a systematic review and meta-analysis. Brain Spine. 2021;1:100263.

Hinton ZW, Wu CJ, Lachiewicz PF, et al. Current trends in revision hip arthroplasty: indications and types of components revised. J Arthroplasty. 2022;37(Suppl):S611-S615.e7.

Inoue D, et al. Comparison of postoperative complications and survivorship of total hip and knee arthroplasty in dialysis and renal transplantation patients. J Arthroplasty. 2020;35(4):971-75.

Inoue D, et al. Outcomes of simultaneous bilateral total hip arthroplasty for 256 selected patients in a single surgeon's practice. Bone Joint J. 2021 Jul;103-B(7 suppl B): 116-121.

Iorio R, et al. [Internet]. What are the absolute and relative contraindications to elective primary total joint arthroplasty (TJA), with respect to surgical site infection (SSI) and periprosthetic joint infection (PJI) risk? [Internet]. Second international consensus meeting on musculoskeletal infection 2018: hip and knee. [accessed 2026 May 11]. Available from:

https://www.icmortho.org/_files/ugd/0540e9_f047ffe9ecc543279bba6b84688bc27e.pdf

Jo S, et al. Clinical outcomes of total hip arthroplasty for displaced femoral neck fractures in patients 80 years of age and older selected by clinical frailty score. Hip Pelvis. 2020 Sep;32(3):148-155.

Johns K, Mabrouk A, Tavarez MM. Slipped capital femoral epiphysis. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 [cited 2022 May]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK538302/>

Jolissaint JE, Kammire MS, Springer BD, et al. An update on the management and optimization of the patient with morbid obesity undergoing hip or knee arthroplasty. Orthop Clin North Am. 2023;54(3):251-257.

Kassarjian A, Isern-Kebschull J, Tomas X, et al. Postoperative hip MR imaging. Magn Reson Imaging Clin N Am. 2022;30(4):673-688.

Keller L, Hogan C, Schocket A. The role of metal patch testing in evaluating patients for metallic

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 26 of 32

prosthetic joint failure. *Ann Allergy Asthma Immunol.* 2021;125(5):542-547.

Kelly D. Congenital anomalies of the lower extremity. In: Azar FM, Canale T, editors. *Campbell's operative orthopaedics*. 14th ed. Philadelphia (PA): Elsevier; 2021. p. 1080-1185.e11.

Kerr DA, Rosenberg AE. Tumors and tumor-like lesions of joints and related structures. In: Firestein GS, Budd RC, editors. *Firestein & Kelley's textbook of rheumatology*. 11th ed. Philadelphia (PA): Elsevier; 2021. p. 2167-2188.e4.

Khakharia S, Jiranek WA. Leg length inequality: prevention/treatment. In: Berry DJ, editor. *Surgery of the hip*. 2nd ed. Philadelphia (PA): Elsevier; 2020. p. 1272-1280.

Khazi ZM, Cates WT, Westermann RW, et al. Arthroscopy versus open arthrotomy for treatment of native hip septic arthritis: an analysis of 30-day complications. *Arthroscopy*. 2020;36(4):1048-1052.

Kim CH, et al. Clinical outcomes following primary hip replacement arthroplasties in patients with solid organ transplantation: A systematic review and meta-analysis. *Hip Pelvis*. 2022 Sep;34(3):127-139.

Kim HKW, Herring JA. Legg-Calvé-Perthes disease. In: Herring JA, editor. *Tachdjian's pediatric orthopaedics*. 6th ed. Philadelphia (PA): Elsevier; 2022. p. 527-580.e10.

Kurmish A. Implant removal in revision hip arthroplasty. In: Berry DJ, editor. *Surgery of the hip*. 2nd ed. Philadelphia (PA): Elsevier; 2020. p. 1050-1061.

Ladd B, Jones KE, Polly DW. Sacroiliac joint pain. In: Steinmetz MP, Berven SH, Benzel EC, editors. *Benzel's spine surgery*. 5th ed. Philadelphia (PA): Elsevier; 2022. p. 1205-1213.e2.

LeBrun DG, Shen TS, Su EP, et al. Hip resurfacing vs total hip arthroplasty in patients younger than 35 years: a comparison of revision rates and patient-reported outcomes. *Arthroplast Today*. 2021;11:229-233.

Leighton JL, Watts E, Gross AE, et al. Bone grafts in hip surgery. In: Berry DJ, editor. *Surgery of the hip*. 2nd ed. Philadelphia (PA): Elsevier; 2020. p. 128-143.

Li J, et al. Comparison of total joint arthroplasty outcomes between renal transplant patients and dialysis patients-a meta-analysis and systematic review. *J Orthop Surg Res*. 2020 Dec 9;15(1):590.

Lieu D, et al. Total hip replacement in haemodialysis or renal transplant patients. *J Orthop Surg (Hong Kong)*. 2014 Dec;22(3):393-8.

Liu Y, et al. Comparing total hip arthroplasty and hemiarthroplasty for the treatment of displaced femoral neck fracture in the active elderly over 75 years old: a systematic review and meta-analysis of randomized control trials. *J Orthop Surg Res*. 2020 Jun 11;15(1):215.

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 27 of 32

Mackenzie CR, Su EP. Surgical treatment of joint diseases. In: Goldman L, Cooney KA, editors. Goldman-Cecil medicine. 27th ed. Philadelphia (PA): Elsevier; 2024. p. 1826-1830.e1.

Major NM, Anderson MW, Dussault R, et al. Hips and pelvis. In: Major NM, Dussault R, editors. Musculoskeletal MRI. 3rd ed. Philadelphia (PA): Elsevier; 2020. p. 347-375.

Mancino F, et al. Metal-on-metal hips: Ten-year clinical and radiographic outcomes of the ADEPT metal-on-metal hip resurfacing and modular total hip arthroplasty. *Journal Clinical Medicine*. 2023 Jan 22;12(3):889.

Martin SD, et al. Hip arthroscopy versus physical therapy for the treatment of symptomatic acetabular labral tears in patients older than 40 Years: a randomized controlled trial. *The American Journal of Sport Medicine*. 2021 APR;49(5):1199-1208.

Mascioli A, Park AL. Infectious arthritis. In: Azar FM, Beaty JH, editors. Campbell's operative orthopaedics. 14th ed. Philadelphia (PA): Elsevier; 2021. p. 485-492.e2.

Mautner K, Pitts J, Street A, et al. Hip injection techniques. In: Frontera WR, Silver JK, Rizzo TD, editors. Atlas of interventional orthopedics procedures. Philadelphia (PA): Elsevier; 2023. p. 323-365.

Mayer SW, Flemig AD, Pearce SS. Hip disorders in the female athlete. In: Frank RM, Matzkin EG, editors. The female athlete. Philadelphia (PA): Elsevier; 2022. p. 139-160.

McGovern RP, Bucci G, Christoforetti JJ, et al. Arthroscopic capsular management of the hip: a comparison of indications for and clinical outcomes of periportal versus interportal capsulotomy. *Arthroscopy*. 2021;37(1):86-94.

Mintz DN, Roberts CC, Weissman BN, et al. ACR Appropriateness Criteria® chronic hip pain. *J Am Coll Radiol*. 2017;14(5 Suppl):S90-S102.

Modest J, Boduch AN, Cruz AI. Slipped capital femoral epiphysis (SCFE). In: Ferri FF, editor. Ferri's clinical advisor 2022. Philadelphia (PA): Elsevier; 2022. p. 1391.e2-1391.e4.

Murugan C, Shetty AP, Rajasekaran S. An unusual case of gluteal pain: case report and review of literature. *J Clin Orthop Trauma*. 2021;20:101487.

Naraghi AM, Mohankumar R, White LM, et al. Bone tumors. *Radiol Clin North Am*. 2022;60(2):205-219.

National Institute for Health and Care Excellence (NICE) [Internet]. Total hip replacement and resurfacing arthroplasty for end-stage arthritis of the hip [TA304]. Original 2014 [accessed 2026 May 11] Available from: Total hip replacement and resurfacing arthroplasty for end-stage arthritis of the hip

Novais E. Treatment options for end-stage hip disease in adolescents: to replace, fuse, or

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 28 of 32

reconstruct? J Pediatr Orthop. 2021;41(Suppl 2):S14-S24.

Parcells BW. Pediatric hip and pelvis. Pediatr Clin North Am. 2020;67(1):139-152.

Patel J, Bozic K. Preoperative planning and templating for revision hip arthroplasty. In: Berry DJ, editor. Surgery of the hip. 2nd ed. Philadelphia (PA): Elsevier; 2020. p. 1035-1049.

Patel KM, Mears SC, Stronach BM, et al. Sepsis and total joint arthroplasty. Orthop Clin North Am. 2022;53(1):13-24.

Patsch JM, Krestan CR. Metabolic and endocrine skeletal disease. In: Adam A, Dixon AK, editors. Grainger & Allison's diagnostic radiology. 7th ed. Philadelphia (PA): Elsevier; 2021. p. 1087-1115.

Peck JB, Sucato DJ. Slipped capital femoral epiphysis. In: Berry DJ, editor. Surgery of the hip. 2nd ed. Philadelphia (PA): Elsevier; 2020. p. 502-517.

Phillips BB, Mihalko MJ. Arthroscopy of the lower extremity. In: Azar FM, Canale T, editors. Campbell's operative orthopaedics. 14th ed. Philadelphia (PA): Elsevier; 2021. p. 2576-2662.e8.

Pisanu F, Andreozzi M, Doria C, et al. Surgical management of hip prosthetic failure in metallosis: a case series and literature review. J Orthop. 2021;28:10-20.

Planka MA, Serino J, DeFroda SF. Greater trochanteric pain syndrome: evaluation and management of a wide spectrum of pathology. SAGE Open Med. 2021;9:1-12.

Popat R, et al. Outcomes of total hip arthroplasty in haemodialysis and renal transplant patients: systematic review. Hip Int. 2021 Mar;31(2):207-214.

Prabowo Y, Saleh I, Junaidi A, et al. Management of synovial chondromatosis of the hip by open arthrotomy debridement only vs total hip replacement: a case report. Int J Surg Case Rep. 2020;74:289-295.

Prince WB. Pediatric musculoskeletal disorders. In: Walls RM, editor. Rosen's emergency medicine: concepts and clinical practice. 10th ed. Philadelphia (PA): Elsevier; 2023. p. 2201-2218.e1.

Rees HW. Management of osteoarthritis of the hip. J Am Acad Orthop Surg. 2020;28(7):e288-e291.

Ren R, et al. Blood metal ion levels after hip resurfacing: a comparison of two different implants. Arthroplasty Today. 2024;30(101555).

Sadaniantz K, Daly L, Tracy J, et al. Prophylactic fasciotomy of the anterolateral thigh donor site in head and neck reconstruction. Am J Otolaryngol. 2022;43(1):103238.

Sahr ME, Miller TT. Pain after hip arthroplasty. Magn Reson Imaging Clin N Am. 2023;31(2):215-238.

Shane AM, Reeves CL, Calaj PM, et al. Soft tissue pathology. Clin Podiatr Med Surg. 2023;40(3):381-395.

Sheffer B. Osteochondrosis or epiphysitis and other miscellaneous affections. In: Azar FM, Beaty JH, editors. Campbell's operative orthopaedics. 14th ed. Philadelphia (PA): Elsevier; 2021. p. 1246-

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 29 of 32

1314.e13.

Shrader MW, et al. Total hip arthroplasty in patients with renal failure: a comparison between transplant and dialysis patients. *J Arthroplasty*. 2006;21(3):324-329.

Smith TO, et al. The clinical and radiological outcomes of hip resurfacing versus total hip arthroplasty: a meta-analysis and systematic review. *Acta Orthop*. 2010;81(6):684-695.

Southgate RD, Vail TP. Resurfacing hip arthroplasty: evolution, design, indications, techniques, and results. In: Berry DJ, editor. *Surgery of the hip*. 2nd ed. Philadelphia (PA): Elsevier; 2020. p. 779-788.

Stanborough RO, Bestic JM, Peterson JJ, et al. Shoulder osteoarthritis. *Radiol Clin North Am*. 2022;60(4):593-603.

Steffner RJ, Spiguel AR, Balach T. Pathologic fractures. In: Browner BD, Jupiter JB, editors. *Skeletal trauma: basic science, management, and reconstruction*. 6th ed. Philadelphia (PA): Elsevier; 2021. p. 613-646.

Strait AV, Ho H, Sershon RA, et al. Outpatient total joint arthroplasty in the “unhealthy”: staying safe using institutional protocols. *J Arthroplasty*. 2025;40(1):34-39.

Taunton MJ, Trousdale RT. Arthrodesis and resection arthroplasty of the hip. In: Berry DJ, editor. *Surgery of the hip*. 2nd ed. Philadelphia (PA): Elsevier; 2020. p. 739-749.

American Academy of Orthopaedic Surgeons [Internet]. Slipped capital femoral epiphysis (SCFE). Rosemont (IL): American Academy of Orthopaedic Surgeons; 2020 [cited 2022 May]. [accessed 2026 Sept 11]. Available from: <https://orthoinfo.aaos.org/en/diseases--conditions/slipped-capital-femoral-epiphysis-scfe>

Thomas WC, Parvataneni HK, Gray CF, et al. Early polyethylene failure in a modern total hip prosthesis: a note of caution. *J Arthroplasty*. 2020;35(5):1297-1302.

Thompson A, Fleischmann KE, Williams K, et al. 2024 AHA/ACC/ACS/ASNC/HRS/SCAI/SCCT/SCMR/SVM guideline for perioperative cardiovascular management for noncardiac surgery: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2024;150(19):e351-e442.

Toy PC, Heck RK. Soft-tissue tumors. In: Azar FM, Beaty JH, editors. *Campbell's operative orthopaedics*. 14th ed. Philadelphia (PA): Elsevier; 2021. p. 1049-1078.e4.

Treacy RB, et al. Birmingham hip resurfacing: a minimum follow-up of ten years. *J Bone Joint Surg Br*. 2011 Jan;93(1):27-33.

United States Food and Drug Administration (FDA) [Internet]. Medical devices/ metal on metal hip implant systems [Updated 2019 Mar 15; accessed 2026 May 11] Available from: <https://www.fda.gov/medical-devices/metal-metal-hip-implants/metal-metal-hip-implant-systems>

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 30 of 32

Upfill-Brown A, Hsiue PP, Stavrakis AI, et al. Instability is the most common indication for revision hip arthroplasty in the United States: national trends from 2012 to 2018. *Arthroplast Today*. 2021;11:88-101.

Usuki KY, Milano MT, Okunieff P, et al. Metastatic disease: bone, spinal cord, brain, liver, and lung. In: Tepper JE, Foote RL, editors. *Gunderson & Tepper's clinical radiation oncology*. 5th ed. Philadelphia (PA): Elsevier; 2021. p. 461-477.e6.

Vahedi H, Yacovelli S, Parvizi J, et al. Surgical treatment of femoroacetabular impingement. *JBJS Rev*. 2021;6(4).

Van de Sande MGH, Van Vollenhoven RF. Evaluation of monoarticular and polyarticular arthritis. In: Firestein GS, Budd RC, Gabriel SE, McInnes IB, O'Dell JR, editors. *Firestein & Kelley's textbook of rheumatology*. 12th ed. Philadelphia (PA): Elsevier. p. 647-661.

Van Vollenhoven RF. Evaluation of monoarticular and polyarticular arthritis. In: Firestein GS, Budd RC, Gabriel SE, McInnes IB, O'Dell JR, editors. *Firestein & Kelley's textbook of rheumatology*. 11th ed. Philadelphia (PA): Elsevier; 2021. p. 663.

Vanaclocha V, Saiz-Sapena N. Percutaneous radiofrequency hip joint denervation. In: Saiz-Sapena N, Vanaclocha V, editors. *Chronic pain: physiopathology and treatment*. Norderstedt (DE): BoD - Books on Demand; 2021. p. 57-71.

Vullo PA, Moreno O, Vita DC. Haemodynamic instability during hip prosthesis removal due to metallosis: an acute chrome/cobalt intoxication? A case report. *Int J Surg Case Rep*. 2022;94:1-4.

Waldman SD. Percutaneous sacroiliac joint fusion. In: Waldman SD, editor. *Atlas of interventional pain management*. 5th ed. Philadelphia (PA): Elsevier; 2021. p. 1146.

Weber AE, Bell JA, Bolia IK. Hip abductor and peritrochanteric space conditions. *Clin Sports Med*. 2021;40(2):311-322.

Wilkin G. Hip arthroscopy for structural hip problems. In: Berry DJ, editor. *Surgery of the hip*. 2nd ed. Philadelphia (PA): Elsevier; 2020. p. 672-686.

Wong BW, Chandrashekar AS, Martin JR, et al. Should we expand outpatient total joint selection criteria? Re-exploring hard stop criteria. *J Arthroplasty*. 2025;40(10):2852-2858.

Za P, et al. Hip resurfacing versus total hip arthroplasty: a systematic review and meta-analysis of randomized clinical trials. *Int Orthop*. 2024 Oct;48(10):2589-2601.

Zavras AG, Fice MP, Colman MW, et al. Indication for proximal femoral replacement is associated with risk of failure. *J Arthroplasty*. 2022;37(5):917-924.

Zhang W, et al. OARSI recommendations for the management of hip and knee osteoarthritis, part I: critical appraisal of existing treatment guidelines and systematic review of current research evidence. *Osteoarthritis Cartilage*. 2007 Sep;15(9):981-1000.

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 31 of 32

Zhang W, et al. OARSI recommendations for the management of hip and knee osteoarthritis, part II: OARSI evidence-based, expert consensus guidelines. Osteoarthritis Cartilage. 2008 Feb;16(2):137-62.

Zhang W, et al. OARSI recommendations for the management of hip and knee osteoarthritis, part III: Changes in evidence following systematic cumulative update of research published through January 2009. Osteoarthritis Cartilage 2010;18(4):476-499.

Zuke WA, et al. Long-term results of the Birmingham Hip Resurfacing Implant in the United States: an updated analysis of a single institution's experience. J Bone Joint Surg Am. 2025 Mar 19;107(6):e19.

SEARCH TERMS

Not Applicable

CENTERS FOR MEDICARE AND MEDICAID SERVICES (CMS)

[Total Joint Arthroplasty \(LCD L36039\)](#) [accessed 2026 September 10].

[Total Joint Arthroplasty \(LCA A57428\)](#) [accessed 2026 September 10].

Total Hip Resurfacing is not specifically addressed in National or Regional Medicare coverage determinations or policies.

PRODUCT DISCLAIMER

- Services are contract dependent; if a product does not cover a service, medical policy criteria do not apply.
- If a commercial product (including an Essential Plan or Child Health Plus product) covers a specific service, medical policy criteria apply to the benefit.
- If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit.
- If a Medicare product (including Medicare HMO-Dual Special Needs Program (DSNP) product) covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.
- If a Medicare HMO-Dual Special Needs Program (DSNP) product DOES NOT cover a specific service, please refer to the Medicaid Product coverage line.

POLICY HISTORY/REVISION

Committee Approval Dates

09/17/26

Medical Policy: Orthopedic Joint Procedures of the Hip for Managed Medicaid (MMC), Health and Recovery Plan (HARP), Child Health Plus (CHP), and Essential Plan (EP) Members

Policy Number: 7.01.121

Page: 32 of 32

Date	Summary of Changes
01/01/2027	Original Effective Date
09/17/26	New Policy