

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	Radiopharmaceuticals for the Treatment of Cancer
Policy Number	6.01.44
Category	Technology Assessment
Original Effective Date	08/21/14
Committee Approval Date	04/16/15, 04/21/16, 04/20/17, 04/19/18, 05/16/19, 04/16/20, 05/21/20, 05/20/21, 07/21/22, 05/18/23, 09/21/23, 09/19/24
Current Effective Date	09/19/24
Archived Date	N/A
Archive Review Date	N/A
Product Disclaimer	<ul style="list-style-type: none"> • Services are contract dependent; if a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan or Child Health Plus product), medical policy criteria apply to the benefit. • If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit. • If a Medicare product (including Medicare HMO-Dual Special Needs Program (DSNP) product) covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit. • If a Medicare HMO-Dual Special Needs Program (DSNP) product DOES NOT cover a specific service, please refer to the Medicaid Product coverage line.

POLICY STATEMENT

- I. Based upon our criteria and the lack of peer-reviewed literature, requests for radiopharmaceutical treatments in excess of their U.S. Food and Drug Administration (FDA) approved regimen, will be considered **investigational**.
- II. Xofigo injection (Radium Ra 223 Dichloride)
 - A. Based upon our criteria and assessment of the peer-reviewed literature, Xofigo has been medically proven to be effective and, therefore, is considered **medically appropriate** when **ALL** of the following criteria are met:
 1. diagnosis of medically or surgically castration-resistant prostate cancer,
 2. presence of symptomatic bone metastases;
 3. no known visceral metastatic disease or bulky regional lymph nodes greater than three (3) cm on imaging performed within the past 30 days.
- III. Pluvicto (Lutetium Lu 177 Vipivotide Tetraxetan)
 - A. Based upon our criteria and assessment of the peer-reviewed literature, Pluvicto has been medically proven to be effective and, therefore, is considered **medically appropriate** when **ALL** of the following criteria are met:
 1. prostate-specific membrane antigen (PSMA)-positive;
 2. metastatic castration-resistant prostate cancer (mCRPC);
 3. have been treated with at least one (1) androgen-receptor pathway inhibitor (i.e., enzalutamide and/or abiraterone);
 4. have been treated with one (1) or two (2) taxane-based regimens;
 5. have at least one PSMA-positive metastatic lesion and no PSMA-negative lesions* on 68Ga-PSMA-11 or F-18 piflufolastat PSMA PET/CT scan.

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer

Policy Number: 6.01.44

Page: 2 of 13

*PSMA negative lesions are defined as metastatic disease that lacks PSMA uptake including bone with soft tissue components ≥ 1.0 cm, lymph nodes ≥ 2.5 cm in short axis, solid organ metastases ≥ 1.0 cm in size.

IV. Strontium-89 Chloride

Based upon our criteria and assessment of the peer-reviewed literature, Strontium-89 Chloride has been medically proven to be effective and, therefore, is considered **medically appropriate** for the treatment of bone pain in patients with confirmed skeletal metastases.

V. Lutathera (lutetium or Lu 177 dotatate)

A. Based upon our criteria and assessment of the peer-reviewed literature, peptide receptor radionuclide therapy using Lutathera has been medically proven to be effective and, therefore, is considered **medically appropriate** when **ALL** of the following requirements are met:

1. Treatment is for **any** of the following:

- a. somatostatin receptor-positive gastroenteropancreatic neuroendocrine tumors (GEP-NETS) of the foregut, midgut and hindgut that are either inoperable or metastatic;
- b. somatostatin receptor positive tumors of the pancreas that are either inoperable metastatic;
- c. somatostatin receptor positive bronchopulmonary or thymic-tumors, which are either inoperable or metastatic; or
- d. pheochromocytomas or paragangliomas;

AND

2. for well-differentiated G1 or G2 neuroendocrine tumors with a Ki-67 $< 20\%$; or well-differentiated G3 neuroendocrine tumors with a Ki-67 $< 55\%$;
3. positive somatostatin receptor scintigraphy with correlative MRI or CT imaging of metastatic measurable disease or 68-Ga-Dotate PET scan positive for metastatic disease*;
4. progression of the disease following treatment with somatostatin-analogs (SSA);
5. the absence of all of the following contraindications:
 - b. serum creatinine: 1.7 mg or greater per deciliter or creatinine clearance of less than 50 ml/minute;
 - c. Hgb. 8.0 g/dl or less; WBC less than 2000/mm³; platelets less than 75,000 mm³.

*In the absence of metastatic disease, documentation should include a surgical or medical consult and the reason for inoperability.

VI. Azedra (iobenguane I 131)

A. Based upon our criteria and assessment of the peer-reviewed literature, Azedra (iobenguane I 131) has been medically proven to be effective and therefore, is considered **medically appropriate** when **ALL** of the following criteria are met:

1. inoperable locally advanced or metastatic pheochromocytoma or paraganglioma requiring systemic treatment;
2. individual is 12 years of age or older;
3. iodine meta-iodobenzylguanidine scan (MIBG, Iobenguane scan) positive;
4. the absence of **all** of the following contraindications:
 - a. creatinine clearance < 30 mL/min;
 - b. platelet count of 80,000/mcL or less;
 - c. absolute neutrophil count of 1,200/mcL or less.

VII. HICON (sodium iodide I 131 solution/ capsules)

A. Based upon our criteria and assessment of the peer-reviewed literature, HICON (sodium iodide I 131 solution/ capsules) have been medically proven to be effective and therefore, are considered **medically appropriate** for the treatment of thyroid carcinoma (i.e., follicular, papillary, Hürthle cell) when **ANY** of the following criteria are met:

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer

Policy Number: 6.01.44

Page: 3 of 13

1. remnant ablation after surgery in T1b/T2 localized disease when there are no other adverse pathologic, laboratory, or imaging features;
2. adjuvant therapy under **any** of the following circumstances:
 - a. gross extrathyroidal extension;
 - b. a primary tumor greater than 4 cm;
 - c. postoperative unstimulated thyroglobulin (Tg) greater than 10ng/mL;
 - d. six or more positive lymph nodes or bulky lymph nodes;
 - e. for follicular or Hürthle cell/Oncocytic, extensive vascular invasion (≥ 4 foci);
3. presence of proven and documented unresectable or metastatic disease based on pathology or pretherapy radioiodine scan.

VIII. Zevalin (ibritumomab tiuxetan)

- A. Based upon our criteria and assessment of the peer-reviewed literature, Zevalin (ibritumomab tiuxetan) has been medically proven to be effective, and therefore, is considered **medically appropriate** for adults when **ALL** of the criteria are met:
1. Treatment is for any of the following:
 - a. relapsed low grade B-cell CD20-positive follicular non-Hodgkin lymphomas (NHLs);
 - b. refractory low grade B-cell CD20-positive follicular NHLs;
 - c. newly diagnosed (consolidation after chemotherapy) low grade B-cell CD20-positive NHLs after at least a partial response (PR) to therapy;
 - d. newly diagnosed (initial treatment) low grade B-cell CD20-positive follicular NHLs for the elderly or infirm when no other option is expected to be tolerated; **or**
 - e. transformed B-cell follicular NHLs that are CD20-positive;

AND

2. bone marrow biopsy demonstrating <25% involvement with lymphoma;

AND

3. the absence of **all of** the following contraindications:
 - a. platelet count of 100,000/mm³ or less;
 - b. neutrophil count (ANC) $\leq 1,500$, bone marrow cellularity < 15%;
 - c. high tumor burden in the bone marrow (lymphoma bone marrow involvement >25%);
 - d. previous radiation to >25% of active marrow sites.

This policy does not address the use of radiotracers for advanced diagnostic procedures. Please refer to CMP #6.01.29 Positron Emission Tomography (PET) Oncologic Applications

Refer to Corporate Medical Policy #11.01.03 Experimental or Investigational Services

POLICY GUIDELINES

- I. The dose regimen of Xofigo is 55 kBq (1.49 mCi) per kg body weight, given in six injections at four (4) week intervals.
- II. The dose regimen of Pluvicto is 7.4 GBq (200 mCi) given by intravenous (IV) injection or infusion every six (6) weeks for up to six (6) doses. The frequency depends on how the cancer responds and how the patient tolerates therapy.
- III. The recommended dose of Strontium-89 Chloride is 148 MBq, 4 mCi, administered IV over one (1) to two (2) minutes or a dose of 1.5-2.2 MBq/kg, 40-60 uCi/kg body weight. Repeated administration of Strontium-89 Chloride should be based on an individual patient's response to therapy, current symptoms, and hematologic status, and are generally not recommended at intervals of less than 90 days.
- IV. The recommended dose of Lutathera is 7.4 GBq (200 mCi) every 8 weeks for a total of four (4) doses.

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer

Policy Number: 6.01.44

Page: 4 of 13

- V. The recommended dose regimen of Azedra is based on body weight. The dosimetric dose for patients weighing greater than 50 kg is 185 to 222MBq (5 or 6 mCi). The dosimetric dose for patients weighing 50 kg or less is 3.7 MBq/kg (0.1 mCi/kg)
- VI. The recommended dose regimen of HICON is based on thyroid gland size and uptake. The dose for the treatment of thyroid carcinoma is 1,110 MBq to 3,700 MBq (30 mCi to 100 mCi) administered orally. The National Comprehensive Cancer Network (NCCN) recommends 30-50mCi for remnant ablation, 75-150mCi for adjuvant therapy.
- VII. The recommended dose regimen of Zevalin is administered in two (2) steps and is based on body weight; 0.4mCi/kg if platelet count is $\geq 150,000$; 0.3 mCi/kg (11.1MBq per kg) if platelet count is 100,000-149,000/mm³.

DESCRIPTION

Radiopharmaceuticals deliver radiation to the cancer cells within their microenvironment, providing a more targeted approach. This is done either by using delivery vehicles that bind preferentially to a specific target or may be taken up by a tumor based on its environment. Radiopharmaceuticals have different emission properties which deliver radiation using either alpha or beta particles. Response to treatment with radiopharmaceuticals may occur after one single injection or up to five injections, and therefore, is quicker than chemotherapy which may occur after many months/cycles. Adverse events from treatment may also be less.

Xofigo

On May 15, 2013, the FDA approved radium-223 marketed under the name Xofigo injection (Bayer HealthCare Pharmaceuticals Inc). Xofigo is an alpha-emitting agent for treatment of patients with symptomatic, bone-metastatic, castration resistant prostate cancer (CRPC). Xofigo has a half-life of 11.4 days, and releases 94% of its energy as alpha-particles with very little beta or gamma-emission. Xofigo mimics calcium and forms complexes with the bone mineral hydroxyapatite at areas of increased bone turnover, such as bone metastases. Alpha-emission consists of particles with high energy and a short range, causing non-repairable breakage of double-strand DNA in adjacent cells, which results in a highly localized cytotoxic effect in the target areas, and causes an anti-tumor effect on bone metastases. The alpha particle range from Xofigo is less than 100 micrometers (less than 10 cell diameters) which limits damage to the surrounding normal tissue and reduces marrow toxicity.

Xofigo is administered intravenously, given once per month for six months, by an appropriately licensed facility, usually in nuclear medicine or radiation therapy departments. Hematologic evaluation of patients must be performed at baseline and prior to every dose of Xofigo. Prior to the initial dose, patients must have absolute neutrophil count (ANC) greater than or equal to $1.5 \times 10^9/L$, platelet count greater than or equal to $100 \times 10^9/L$, and hemoglobin greater than or equal to 10 g/dL. Before subsequent administrations of Xofigo, the ANC should be greater than or equal to $1 \times 10^9/L$ and the platelet count greater than or equal to $50 \times 10^9/L$. Xofigo should be discontinued if a delay of six to eight weeks does not result in the return of blood counts to these levels. Non-hematologic side effects are generally mild, and include nausea, diarrhea, and vomiting. These symptoms are likely related to the fact that Xofigo is predominantly eliminated by fecal excretion. Whenever possible, patients should use a toilet and the toilet should be flushed several times after each use. Clothing soiled with Xofigo or patient fecal matter or urine should be washed promptly and separately from other clothing. Caregivers should use universal precautions, such as washing hands, wearing gloves and barrier gowns when handling patients' bodily fluids to avoid contamination.

Pluvicto

Formally known as ¹⁷⁷Lu-PSMA-617, Pluvicto (Novartis) was approved by the FDA on March 23, 2022 for the treatment of adult patients with prostate-specific membrane antigen (PSMA)-positive metastatic castration-resistant prostate cancer who have been treated with androgen receptor (AR) pathway inhibition and taxane-based chemotherapy.

Pluvicto is the first targeted radioligand therapy. The half-life is 6.647 days by emitting beta-particle radiation. Pluvicto emits beta-particle radiation selectively to PSMA-positive cells surrounding microenvironment. In early-phase studies in

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer

Policy Number: 6.01.44

Page: 5 of 13

previously treated patients with metastatic castration resistant prostate cancer, Pluvicto has shown biochemical and radiographic response rates, reduced pain, and low toxicity.

Strontium-89 Chloride

On July 11, 2013, the FDA approved Strontium-89 chloride for relief of bone pain in patients with painful skeletal metastases. Strontium-89 Chloride acts similarly to calcium, localizing in bone mineral and is absorbed in sites of osteogenesis, especially at sites of metastases, compared to normal healthy bone. Strontium-89 Chloride is a beta-emitter with a maximum range in tissue of ~ 8 mm and has a half-life of 50.5 days. The recommended dose of Strontium-89 Chloride is 148 MBq, 4 mCi, administered IV over one to two minutes or a dose of 1.5- 2.3 MBq/kg, 40-60 uCi/kg body weight. Repeated administration of Strontium-89 Chloride should be based on an individual patient's response to therapy, current symptoms, and hematologic status, and are generally not recommended at intervals of less than 90 days. Bone marrow toxicity in the white blood cells and platelets is variable from patient to patient and must be monitored following administration of Strontium-89 Chloride. Relief of pain may not occur for seven to 20 days post injection and some patients have experienced a transient increase in pain at 36 to 72 hours after injection. Strontium-89 Chloride is excreted by the kidney and should be administered with caution in patients with renal dysfunction.

Lutathera (lutetium or Lu 177 dotatate)

Lutathera (lutetium or Lu 177 dotatate) (Novartis, formerly Advanced Accelerator Applications) was approved by the FDA on January 26th, 2018. It is also commonly referred to as "Lu 177". It is classified as a peptide receptor radionuclide. It is a radiolabeled-somatostatin analogue that binds to somatostatin receptor expressing cells, including malignant somatostatin receptor-positive tumors such as neuroendocrine tumors. It is then internalized and beta particle emission from Lutathera induces cellular damage by formation of free radicals in somatostatin receptor-positive cells and adjacent cells. It is considered as an alternative treatment option to first-line somatostatin analogues.

Azedra (Iobenguane I 131)

Azedra (Iobenguane I 131) (Progenics Pharmaceuticals, Inc.) was approved by the FDA on July 30th, 2018 and is a radioactive therapeutic agent with a structure similar to norepinephrine. Due to its structure, it is taken up by the norepinephrine transporter where it accumulates in adrenergically innervated tissues including pheochromocytoma and paraganglioma cells. The beta and gamma radiation resulting from the radioactive decay causes an anti-tumor effect. It is indicated for the treatment of adult and pediatric patients 12 years of age and older with iobenguane scan positive, unresectable, locally advanced or metastatic pheochromocytoma or paraganglioma who require systemic therapy.

HICON (Sodium iodide I 131)

The primary treatment of choice for differentiated thyroid carcinoma is surgery, followed by radioactive iodine (RAI) ablation for select patients. HICON (Sodium iodide I 131) also referred to as ¹³¹I, is a radioactive therapeutic agent, when given orally, (via capsule or solution) is naturally taken up through the blood by the thyroid, causing decay via beta emission and associated gamma emission. RAI has been in use since the 1940's, but the high concentration version of the drug received FDA approval on January 24th, 2003 (DraxImage). Serum thyroglobulin (Tg) levels can indicate the existence of distant metastases. RAI is given after total thyroidectomy to eliminate the normal thyroid remnant, ensure that Tg levels are undetectable after surgery, to reduce the risk of recurrence (adjuvant treatment), and to treat persistent or recurrent disease (treatment of known disease). Tumors that do not take up iodine, i.e., anaplastic (undifferentiated) and medullary thyroid carcinomas cannot be treated with RAI.

Zevalin (ibritumomab tiuxetan)

Zevalin (ibritumomab tiuxetan) (Biogen Idec) is a combination of a CD20-directed monoclonal antibody and the radioisotope yttrium-90. It has also been referred to as ⁹⁰Y ibritumomab tiuxetan. Zevalin was FDA approved on February 19, 2002 for the treatment of patients with relapsed or refractory low-grade, follicular, or transformed B-cell Non-Hodgkin lymphomas (NHL), including patients with Rituximab refractory follicular NHL. The regimen includes the unlabeled antibody rituximab, which is given prior to Zevalin to saturate CD20 binding sites on B-cells in the peripheral

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer

Policy Number: 6.01.44

Page: 6 of 13

blood and spleen, to improve Zevalin's biodistribution. Zevalin is seen as an effective treatment alternative for eligible patients but is not commonly used because the administration is seen as highly complex.

RATIONALE

The National Comprehensive Cancer Network (NCCN) Guidelines for Prostate Cancer (v4.2024) principles of radiopharmaceutical therapy states that Radium-223 is an alpha-emitting radiopharmaceutical that has been shown to extend survival in patients who have castration-resistant prostate cancer (CRPC) with symptomatic bone metastases, but no visceral metastases. Radium-223 alone has not been shown to extend survival in patients who have visceral metastases or bulky nodal disease greater than three to four cm. Radium-223 differs from beta-emitting agents, such as samarium-153 and strontium-89, which are palliative and have no survival advantage. Radium-223 causes double-strand DNA breaks and has a short radius of activity. Grade 3-4 hematologic toxicity (2 percent neutropenia, 3 percent thrombocytopenia, 6 percent anemia) occurs at a low frequency. At the present time, except on a clinical trial, Radium-223 is not intended to be used in combination with chemotherapy due to the potential for additive myelosuppression. Concomitant use of denosumab or zoledronic acid does not interfere with the beneficial effects of Radium-223 on survival.

The Alphasradin in Symptomatic Prostate Cancer Patients (ALSYMPCA) trial (Parker, et al. 2013) was a phase 3, randomized, double-blind, placebo-controlled study that randomized 921 patients with symptomatic bone-metastatic CRPC to six injections every week of either radium-223(50 kBq/kg) or placebo. Patients were symptomatic with two or more bone metastases, without visceral metastases and had received docetaxel or were ineligible for docetaxel treatment. Median overall survival in the Xofigo arm was 14.9 months compared to 11.2 months in the placebo arm. Median time-to-first skeletal related event was significantly improved in the treatment arm (13.6 months) compared to placebo (8.4 months). Time-to-alkaline-phosphatase-progression and time-to-PSA-progression was also improved in the treatment group. More adverse events were observed in the Xofigo group, with discontinuation of treatment due to adverse events occurring in 13 percent of the patients in the Xofigo and 20 percent of the patients in the placebo arm. The significantly improved overall survival in the treatment group met the predetermined boundary for discontinuing the study early and the trial was terminated due to evidence of significant treatment benefit of Xofigo.

The ERA 223 (Smith M, et al. 2019) was a randomized, double-blind, placebo-controlled phase 3 trial. It included 165 oncology and urology centers in 19 countries, which assessed the efficacy and safety of combination therapy with abiraterone acetate plus prednisone or prednisolone and Xofigo in patients with asymptomatic or mildly symptomatic, chemotherapy-naïve, castration-resistant prostate cancer and bone metastases. A total of 806 patients with a minimum of two bone metastases and no known visceral or brain metastases were randomized 1:1 to either Xofigo combination therapy or placebo combination therapy. The primary end-point was occurrence of a symptomatic skeletal event, defined as use of external beam radiotherapy to relieve skeletal symptoms, a new symptomatic pathological bone fracture, spinal cord compression, or tumor-related orthopedic surgical intervention. The study showed that the combination of abiraterone acetate plus prednisone or prednisolone and Xofigo did not improve symptomatic skeletal event-free survival or overall survival and was associated with an increased frequency of fractures. Based on the results of this study, the manufacturer, Bayer HealthCare Pharmaceuticals Inc, does not recommend Xofigo for use in combination with abiraterone acetate plus prednisone/prednisolone outside of clinical trials.

In early-phase studies in previously treated patients with metastatic castration resistant prostate cancer, Lutetium Lu 177 vipivotide tetraxetan (Pluvicto) has shown biochemical and radiographic response rates, reduced pain, and low toxicity.

The VISION trial, an international multi-center phase 3 trial compared treatment with Pluvicto and standard care to standard care alone in patients with metastatic castration-resistant prostate cancer previously treated with at least one androgen-receptor-pathway inhibitor and one or two taxane regimens (Sartor et al., 2021). A total of 891 patients with a PSMA-positive gallium-68-labeled PSMA PET scan and who had progressed on androgen-receptor-pathway inhibitors and taxane therapy were randomized 2:1 to Pluvicto and standard care or standard care alone. Imaging-based progression free survival (PFS), overall survival, objective response, disease control, and time to symptomatic skeletal events were evaluated. Both PFS and overall survival were longer in the Pluvicto and standard care group (median 8.8 months and 15.3 months, respectively) compared to standard care alone (3.4 months and 11.3 months, respectively). Pluvicto and standard care group experienced more grade 3 and 4 adverse events (52.7%) than the standard care group (38.0%) but

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer

Policy Number: 6.01.44

Page: 7 of 13

quality of life was not adversely affected. The authors concluded that Pluvicto along with standard care prolongs PFS and overall survival when compared to standard care alone in patients with metastatic castration – resistant prostate cancer.

The NCCN Guidelines for Bone Cancer (v2.2024) state that samarium-153 ethylene diamine tetramethylene phosphonate (Sm-153-EDTMP) is a beta-particle-emitting, bone-seeking radiopharmaceutical, and has been evaluated in patients with locally recurrent or metastatic osteosarcoma or skeletal metastases. Studies have reported that Sm-153-EDTMP with peripheral blood progenitor cell support had low non-hematologic toxicity and provide pain palliation for patients with osteosarcoma local recurrence or osteoblastic bone metastases. Results of dose-finding studies have also demonstrated that Sm-153-EDTMP can be effective in the treatment of patients with high-risk osteosarcoma.

The FDA approval for the use of Lutathera is based on the results of two published studies. The NETTER 1 study (Strosberg et al. 2021) compared treatment with Lutathera to octreotide in patients with inoperable, progressive somatostatin receptor-positive midgut carcinoid tumors. Eligibility included a Ki67 index of 20% or lower, OctreoScan uptake greater than or equal to that of the normal liver, creatinine clearance of 50 mL/min or greater, no prior treatment with PRRT, and no prior external radiation therapy to more than 25% of the bone marrow. The primary outcome was progression free survival (PFS). A total of 229 patients were randomized to Lutathera 200 mCi for four infusions every eight weeks concurrently with long-acting octreotide (30 mg) or high-dose octreotide alone (60 mg). Baseline characteristics were balanced between the groups. It was noted that 74% of patients had an ileal primary and 96% had metastatic disease in the liver. At the data-cutoff date for the primary analysis, PFS at 20 months was 65.2% in the Lutathera arm vs 10.8% in the control group. The response rate was 18% in the Lutathera group vs 3% in the control group. In an updated analysis, progressive disease was seen in 23% of the Lutathera group and 69% of the control group. Median progression free survival was not reached for the experimental group and was 8.5 months for the control group. Median overall survival was also not reached in the experimental group but was 27.4 months in the control arm.

Singh and colleagues (2024) published the phase 3 NETTER-2 trial, assessing the use of ¹⁷⁷Lu-Dotatate in a first-line metastatic setting. Participants (n=226) with newly diagnosed higher grade 2 and grade 3 somatostatin receptor-positive, advanced gastroenteropancreatic neuroendocrine tumors were randomized 2:1. The ¹⁷⁷Lu-Dotatate group received 4 cycles of IV [¹⁷⁷Lu]Lu-DOTA-TATE plus IM octreotide 30mg long-acting repeatable (LAR) followed by octreotide 30 mg LAR every 4 weeks. The control group was given high-dose octreotide 60mg LAR every 4 weeks. Tumor assessments were conducted at baseline, 16 weeks, 24 weeks, and then every 12 weeks until disease progression or death. The primary end point was progression free survival. The ¹⁷⁷Lu-Dotatate group outperformed the control group demonstrating a progression free survival of 22.8 month vs. 8.5 months for the control group. Authors concluded that first line ¹⁷⁷Lu-Dotatate plus octreotide LAR should be considered a new standard of care in first-line therapy for patients with grade 2 or 3 advanced gastroenteropancreatic neuroendocrine tumors.

The MIP-1B12B study (NCT00874614) was a multicenter, open-label, single-arm prospective study of the radiopharmaceutical Azedra, that included 74 intention-to-treat participants, 12 years of age or older with a diagnosis of pheochromocytoma or paraganglioma. To be eligible, participants had to be appropriate for curative treatment and had failed a prior treatment. The primary outcome was the number of patients with 50% or greater reduction in antihypertensive medication(s) lasting for at least six months after administering the last therapeutic dose of Azedra. The primary endpoint was met by 25% of patients (95% CI, 16.2% to 36.5%). Antitumor activity was demonstrated with 22.1% (95% CI, 13.6% to 32.7%) of patients maintaining a response for at least 6 months. Given the severity and rarity of the condition and its association with a high degree of morbidity and mortality, the results of the study represent a clinically meaningful outcome.

In 2019, the American Thyroid Association (ATA), the European Association of Nuclear Medicine and the Society of Nuclear Medicine and Molecular Imaging, and the European Thyroid Association issued a joint statement for the use of ¹³¹I therapy for the treatment of differentiated thyroid cancer. The societies defined terminology that should be used to communicate the goals of ¹³¹I therapy. Adapted from the joint statement:

	¹³¹I Therapy
--	--------------------------------

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer

Policy Number: 6.01.44

Page: 8 of 13

Goal	Remnant Ablation	Adjuvant Treatment	Treatment of Known Disease
Initial staging	✓	✓	✓
Facilitate follow-up	✓	✓	✓
Improve disease-specific survival	-	✓	✓
Decrease recurrence	-	✓	-
Improve progression-free survival	-	✓	✓
Curative intent	-	✓	✓
Palliative intent	-	-	✓

The categories for thyroid cancer staging by the American Thyroid Association are low, intermediate, and high risk, which provides risk for recurrence. Preoperative staging and risk of recurrence determine the need for radiotherapy after surgery and following surveillance. The evidence for the use of ¹³¹I in the post operative setting of differentiated thyroid carcinoma is limited to retrospective studies. Although numerous have been published, findings are inconsistent regarding improvement in outcomes. The joint consensus states that, “Even though most guidelines make recommendations with regard to the postoperative use of ¹³¹I based primarily on staging systems that predict risk of recurrence or disease specific mortality, the actual goal of ¹³¹I therapy can only be determined once the postoperative disease status has been assessed. Regardless of initial risk stratification, patients with biochemical, structural, or functional evidence of persistent disease can only be candidates for “treatment of known disease.” Patients demonstrating no histological, biochemical, or imaging evidence of persistent disease after appropriate initial surgery may be candidates for observation, remnant ablation, or adjuvant treatment.”

NCCN, in the v.3.2024 Guidelines for Thyroid Carcinoma, recommends postoperative RAI when a number of clinical factors predict a significant risk of recurrence, distant metastases, or disease-specific mortality, including carcinoma type, size of primary tumor, post operative Tg level, vascular invasion, presence of metastases, etc. RAI is not typically indicated for patients considered to have a low risk of recurrence, or after lobectomy, or for patients that have metastatic disease that is not amenable to RAI therapy, meaning iodine refractory disease. NCCN states that RAI may be used for patients without gross residual disease, but that data is conflicting.

Witzig et al., in a 2007 post-hoc analysis, aimed to assess the durability of four clinical trials—a Phase I-II dose-finding trial in patients with indolent and aggressive NHL, a Phase II trial of reduced-dose ⁹⁰Y ibritumomab tiuxetan in patients with mild thrombocytopenia, a randomized Phase III trial that compared the ibritumomab tiuxetan regimen with rituximab mono-therapy, and a Phase III trial in patients with rituximab-refractory NHL. The trials represented a total of 211 patients from 30 centers across the United States. The analysis revealed that 37% of patients treated with ⁹⁰Y ibritumomab tiuxetan had a long-term response to the treatment (defined as time to progression of 12 months or longer). With a median follow-up of 53.5 months, the median duration of response was 28.1 months. The median overall survival in the 211 patients was 49.3 months. Overall survival was 52 months or longer in 82% of those individuals with a long-term response, compared to 34.6 months in patients without a long term response. The estimated overall survival at 5 years was 53% for all patients treated with ⁹⁰Y ibritumomab tiuxetan.

CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member’s subscriber contract.
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer**Policy Number: 6.01.44****Page: 9 of 13**

- *Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN).*

CPT Codes

Code	Description
77750	Infusion or instillation of radioelement solution (includes 3-month follow-up care)
78800	Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); planar, single area (e.g., head, neck, chest, pelvis), single day imaging.
78801	Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); planar, 2 or more areas (e.g., abdomen and pelvis, head and chest), 1 or more days imaging or single area imaging over 2 or more days.
78802	Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); planar, whole body, single day imaging.
78803	Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); tomographic (SPECT), single area (e.g., head, neck, chest, pelvis), single day imaging.
78804	Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); planar, whole body, requiring 2 or more days imaging.
79101	Radiopharmaceutical therapy, by intravenous administration
79005	Radiopharmaceutical therapy, by oral administration
79403	Radiopharmaceutical therapy, radiolabeled monoclonal antibody by intravenous infusion (<i>effective 01/01/04</i>).
96374	Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); intravenous push, single or initial substance/drug.
96375	Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); each additional sequential intravenous push of a new substance/drug (list separately in addition to code for primary procedure).
96409	Chemotherapy administration,; intravenous, push technique, single or initial substance/drug.

Copyright © 2024 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
A4641	Radiopharmaceutical, diagnostic, not otherwise classified
A9513	Lutetium lu 177, dotatate injection, therapeutic, 1mCi (Lutathera)
A9517	Iodine I-131 sodium iodide capsule(s), therapeutic, per mCi (HICON)
A9530	Iodine I-131 sodium iodide solution, therapeutic, per mCi (HICON)

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer**Policy Number: 6.01.44****Page: 10 of 13**

Code	Description
A9543	Yttrium Y-90 ibritumomab tiuxetan, therapeutic, per treatment dose, up to 40 mCi (Zevalin)
A9590	Iodine I-131, Iobenguane, 1mCi (Azedra)
A9600	Strontium sr-89 chloride, therapeutic, per mCi
A9606	Radium RA-223 dichloride, therapeutic, per UCI (Xofigo)
A9607	Lutetium lu 177 vipivotide tetraxetan, therapeutic, 1 millicurie (Pluvicto)
C9399	Unclassified drugs or biologicals

ICD10 Codes

Code	Description
C61	Malignant neoplasm of prostate
C7A.010- C7A.019	Malignant carcinoid tumors of the small intestine code range
C7A.020- C7A.025	Malignant carcinoid tumors of the appendix, large intestine, and rectum code range
C71.029	Malignant carcinoid tumor of the large intestine, unspecified portion
C7A.092	Malignant carcinoid tumor of the stomach
C7A.094- C7A.096	Malignant carcinoid tumor of the foregut, midgut, and hindgut, unspecified code range
C7B.00-C7B.09	Secondary neuroendocrine tumor code range
C7B.8	Other secondary neuroendocrine tumors
C79.51	Secondary malignant neoplasm of bone
C79.52	Secondary malignant neoplasm of bone marrow
D40.0	Neoplasm of uncertain behavior of prostate
V58.0	Radiotherapy
Z85.46	Personal history of malignant neoplasm of prostate

REFERENCES

*A Study Evaluating Ultratrace Iobenguane I131 in Patients with Malignant Relapsed/Refractory Pheochromocytoma/Paraganglioma. ClinicalTrials.gov identifier: NCT00874614. Last updated February 2, 2020. [<https://clinicaltrials.gov/study/NCT00874614>] accessed 08/01/24.

*Barber TW, et al. The potential for induction peptide receptor chemoradionuclide therapy to render inoperable pancreatic and duodenal neuroendocrine tumours resectable. *Eur J Surg Oncol* 2012 Jan;38(1):64-71.

*Bodei L, et al. Yttrium-labeled peptides for therapy of NET. *Eur J Nucl Med Mol Imaging* 2012 Feb;39(Suppl 1):S93-102.

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer

Policy Number: 6.01.44

Page: 11 of 13

Cheng S, et al. Population-based analysis of the use of radium-223 for bone-metastatic castration-resistant prostate cancer in Ontario, and of factors associated with treatment completion and outcome. Cancer Manag Res 2019 Nov; 11:9307-9319.

*Chin RI, et al. Radiopharmaceuticals for neuroendocrine tumors. Semin Radiat Oncol 2021 Jan;31(1):60-70.

*Cooper DS, et al. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. Thyroid 2009;19(11):1167-214.

Den RB, et al. Ra-223 treatment for bone metastases in castrate-resistant prostate cancer: practical management issues for patient selection. Am J Clin Oncol 2019 Apr;42(4):399-406.

*Hamiditabar M, et al. Peptide receptor radionuclide therapy with 177Lu-Octreotate in patients with somatostatin receptor expressing neuroendocrine tumors: six years' assessment. Clin Nucl Med 2017 June;42(6):436-443.

*Harrison MR, et al. Radium-223 chloride: a potential new treatment for castration-resistant prostate cancer patients with metastatic bone disease. Cancer Manag Res 2013;5:1-14.

Heidegger I, et al. Biomarkers to personalize treatment with 177Lu-PSMA-617 in men with metastatic castration-resistant prostate cancer- a state of the art review. Ther Adv Med Oncol 2022; 14: 17588359221081922.

*Heidenreich A, et al. EAU guidelines on prostate cancer. Part II: treatment of advanced, relapsing, and castration-resistant prostate cancer. Eur Urol 2014 Feb;65(2):467-79.

Heinrich D, et al. The contemporary use of radium-223 in metastatic castration-resistant prostate cancer. Clin Genitourin Cancer 2018 Feb;16(1):e223-e231.

*Iyer R, et al. Recent advances in the management of gastroenteropancreatic neuroendocrine tumors: Insights from the 2017 ASCO Gastrointestinal Cancers Symposium. Clin Adv Hematol Oncol 2017 April;15(4 suppl 4):1-24.

Kapoor R, et al. Cancer induced bone pain: current management and future perspectives. Med Oncol 2021 Sep 28;38(11):134.

*Khanna G, et al. Utility of radiolabeled somatostatin receptor analogues for staging/restaging and treatment of somatostatin receptor-positive pediatric tumors. Oncologist 2008 Apr;13(4):382-9.

*Kong G, et al. High clinical and morphologic response using 90Y-DOTA-octreotate sequenced with 177Lu-DOTA-octreotate induction peptide receptor chemoradionuclide therapy (PRCRT) for bulky neuroendocrine tumors. Eur J Nucl Med Mol Imaging 2017 March;44(3):476-489.

*Kong G, et al. Efficacy of peptide receptor radionuclide therapy for functional metastatic paraganglioma and pheochromocytoma. J Clin Endocrinol Metab 2017 Sept 1;102(9):3278-3287.

*Kunikowska J, et al. Long-term results and tolerability of tandem peptide receptor radionuclide therapy with 90Y/177Lu-DOTATATE in neuroendocrine tumors with respect to the primary location: a 10-year study. Ann Nucl Med 2017 June;31(5):347-356.

*Mazzaferri EL, et al. Long-term outcome of patients with differentiated thyroid carcinoma: effect of therapy. Endocrine Practice 2000 Nov/Dec;6(6):469-476.

Murray I and Du Y. Systemic radiotherapy of bone metastases with radionuclides. Clin Oncol (R Coll Radiol) 2021 Feb;33(2):98-105.

National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. B-cell Lymphomas. V2.2024[https://www.nccn.org/professionals/physician_gls/pdf/b-cell.pdf] accessed 08/01/24.

National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Bone Cancer. V2.2024 [http://www.nccn.org/professionals/physician_gls/pdf/bone.pdf] accessed 08/01/24.

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer

Policy Number: 6.01.44

Page: 12 of 13

National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Thyroid Carcinoma. V3.2024[https://www.nccn.org/professionals/physician_gls/pdf/thyroid.pdf] accessed 08/01/24.

National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Neuroendocrine and Adrenal Tumors. V.2.2024. [https://www.nccn.org/professionals/physician_gls/PDF/neuroendocrine.pdf] accessed 08/01/24.

National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Prostate Cancer. V.1.2024. [http://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf] accessed 08/01/24.

*Nilsson S, et al. Two-year survival follow-up of the randomized, double-blind, placebo-controlled phase II study of radium-223 chloride in patients with castration-resistant prostate cancer and bone metastases. Clin Genitourin Cancer 2013 Mar;11(1):20-6.

O'Sullivan JM, et al. Radium-223 within the evolving treatment options for metastatic castration-resistant prostate cancer: recommendations from a European Expert Working Group. Eur Urol Oncol 2020 Aug; 3(4):455-463.

*Pandit-Taskar N, et al. Bone-seeking radiopharmaceuticals for treatment of osseous metastases, Part 1: α therapy with 223Ra-dichloride. J Nucl Med 2014 Feb;55(2):268-74.

*Parghane RV, et al. Clinical response profile of metastatic/advanced pulmonary neuroendocrine tumors to peptide receptor radionuclide therapy with 177Lu-DOTATATE. Clin Nucl Med 2017 June;42(6):428-435.

*Parker C, et al. Alpha emitter radium-223 and survival in metastatic prostate cancer. NEJM 2013;369(3):213-23.

*Parker C, et al. A randomized, double-blind, dose-finding, multicenter, phase 2 study of radium chloride (ra 223) in patients with bone metastases and castration-resistance prostate cancer. Eur Urol 2013 Feb;63(2):189-97.

*Sartor O, et al. Effect of radium-223 dichloride on symptomatic skeletal events in patients with castration-resistant prostate cancer and bone metastases: results from a phase 3, double-blind randomized trial. Lancet Oncol 2014 Jun;15(7):738-46.

*Sartor O, et al. Lutetium-177-PSMA-617 for metastatic castration-resistant prostate cancer. N Engl J Med 2021 Sep 16;385(12):1091-1103.

*Sartor O, et al. Re-treatment with radium-223: 2-year follow-up from an international, open-label, phase I/II study in patients with castration-resistant prostate cancer and bone metastases. Prostate 2019 Oct;79(14):1683-1691.

*Savelli G, et al. Final results of a phase 2A study for the treatment of metastatic neuroendocrine tumors with a fixed activity of (90) Y-DOTA-D-Phe1-Tyr3 octreotide. Cancer 2012 Jun 1;118(11):2915-24.

*Singh S, et al. [^{177}Lu]Lu-DOTA-TATE plus long-acting octreotide versus high-dose long-acting octreotide for the treatment of newly diagnosed, advanced grade 2–3, well-differentiated, gastroenteropancreatic neuroendocrine tumours (NETTER-2): an open-label, randomised, phase 3 study. Lancet 2024;403:2807-2817.

*Smith M, et al. Addition of Radium-223 to abiraterone acetate and prednisone or prednisolone in patients with castration-resistant prostate cancer and bone metastases (ERA 223): a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet Oncol 2019;20(3):408-419.

*Strosberg J, et al. NETTER-1 phase III in patients with midgut neuroendocrine tumors treated with 177Lu-Dotatate: efficacy and safety results. J Nucl Med 2016 May 1;57(Suppl 2):269.

*Strosberg J, et al. Phase 3 trial of 177Lu-Dotatate for midgut neuroendocrine tumors. N Engl J Med 2017 Jan 12;376(2):125-135.

*Strosberg, Jonathan R, et al. ^{177}Lu -Dotatate plus long-acting octreotide versus high-dose long-acting octreotide in patients with midgut neuroendocrine tumours (NETTER-1): final overall survival and long-term safety results from an open-label, randomized, controlled, phase 3 trial. Lancet Oncol 2021 Dec;22(12):1752-1763.

Medical Policy: Radiopharmaceuticals for the Treatment of Cancer

Policy Number: 6.01.44

Page: 13 of 13

*The U.S. Food and Drug Administration. Drugs @FDA: FDA-approved drugs, Azedra. [https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/209607s000lbl.pdf] accessed 08/05/24.

*The U.S. Food and Drug Administration. Drugs @FDA: FDA-approved drugs, HICON. [https://www.accessdata.fda.gov/drugsatfda_docs/label/2003/21305_Sodium-Iodide_lbl.pdf] accessed 08/05/24.

*The U.S. Food and Drug Administration. Drugs @FDA: FDA-approved drugs, Lutathera. [https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/208700s000lbl.pdf] accessed 08/05/24.

*The U.S. Food and Drug Administration. Drugs @FDA: FDA-approved drugs, Pluvicto. [https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/215833s000lbl.pdf] accessed 08/05/24.

*The U.S. Food and Drug Administration. Drugs @FDA: FDA-approved drugs, Xofigo [https://www.accessdata.fda.gov/drugsatfda_docs/label/2013/203971lbl.pdf] accessed 08/05/24.

*The U.S. Food and Drug Administration. Drugs @FDA: FDA-approved drugs, Zevalin. [https://www.accessdata.fda.gov/drugsatfda_docs/label/2002/ibriide021902LB.pdf] accessed 08/05/24.

Tsukamoto S, et al. Current overview of treatment for metastatic bone disease. Curr Oncol 2021 Aug 29;28(5):3347-3372.

Ueno NT, et al. Phase II study of radium-223 dichloride combined with hormonal therapy for hormone receptor-positive, bone-dominant metastatic breast cancer. Cancer Med 2020 Feb;9(3):1025-1032.

Van der Doelen, et al. Patient selection for Radium-223 therapy in patients with bone metastatic castration resistant prostate cancer: new recommendations and future perspectives. Clin Genitourin Cancer 2019 Apr;17(2):79-87.

*Wtzig TE, et al. Long-term responses in patients with recurring or refractory B-cell non-Hodgkin lymphoma treated with Yttrium 90 Ibritumomab Tiuxetan. Cancer 2007 May 1;109(9):1804-1810.

*Key Article

KEY WORDS

XoFigo, Ra-223, radium-223, Radium Ra 223 Dichloride, Lutetium Lu 177 Vipivotide Tetraxetan, Pluvicto, Strontium-89 Chloride, radiopharmaceutical, Strontium, radiotherapeutic, Lutathera, Peptide receptor radionuclide therapy, PRRT, PRRNT, Receptor-mediated radiotherapy, Radiolabeled nuclide therapy, somatostatin analog, ⁹⁰Y-DOTATOC, ¹⁷⁷Lu-DOTA0, Tyr3, ⁹⁰Y-DOTA0.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

Based on our review, radiopharmaceuticals for the treatment of cancer are not addressed in National or Regional Medicare coverage determinations or policies.